arXiv:1507.00719v2 [math.PR] 27 Feb 2016

Liouville quantum gravity and the Brownian map I: The QLE(8/3, 0) metric

Jason Miller and Scott Sheffield

Abstract

Liouville quantum gravity (LQG) and the Brownian map (TBM) are two distinct models of measure-endowed random surfaces. LQG is defined in terms of a real parameter γ, and it has long been believed that when $\gamma = \sqrt{8/3}$, the LQG sphere should be equivalent (in some sense) to TBM. However, the LQG sphere comes equipped with a conformal structure, and TBM comes equipped with a metric space structure, and endowing either one with the *other's* structure has been an open problem for some time.

This paper is the first in a three-part series that unifies LQG and TBM by endowing each object with the other's structure and showing that the resulting laws agree. The present work uses a form of the quantum Loewner evolution (QLE) to construct a metric on a dense subset of a $\sqrt{8/3}$-LQG sphere and to establish certain facts about the law of this metric, which are in agreement with similar facts known for TBM. The subsequent papers will show that this metric extends uniquely and continuously to the entire $\sqrt{8/3}$-LQG surface and that the resulting measure-endowed metric space is TBM.

Contents

1 Introduction **3**
- 1.1 Overview . . . 3
- 1.2 Further observations and sequel overview . . . 10
- 1.3 Prequel overview . . . 11
- 1.4 Outline . . . 14

2 Preliminaries **15**
- 2.1 Continuous state branching processes . . . 15
- 2.2 Quantum surfaces . . . 16
- 2.3 Conformal removability . . . 17

3 Eden model and percolation interface **18**

4 Infinite measures on quantum spheres **22**
- 4.1 Lévy excursion description of doubly-marked quantum spheres . . . 22
- 4.2 Weighted measures . . . 24
- 4.3 Continuum scaling exponents . . . 25

5 Meeting in the middle **26**

6 Quantum natural time $\mathrm{QLE}(8/3, 0)$ **38**
- 6.1 Approximations to $\mathrm{QLE}(8/3, 0)$ . . . 39
- 6.2 Subsequential limits . . . 41
- 6.3 Time parameterizations . . . 45

7 Symmetry **46**
- 7.1 Conditional independence in the limit . . . 46
- 7.2 Reshuffling a pair of SLE_6's . . . 51

8 Metric construction **62**
- 8.1 Setup and notation . . . 63
- 8.2 Coupling explorations and the metric property . . . 63

References **67**

Acknowledgments.

We have benefited from conversations about this work with many people, a partial list of whom includes Omer Angel, Itai Benjamini, Nicolas Curien, Hugo Duminil-Copin, Amir Dembo, Bertrand Duplantier, Ewain Gwynne, Nina Holden, Jean-François Le Gall, Gregory Miermont, Rémi Rhodes, Steffen Rohde, Oded Schramm, Stanislav Smirnov, Xin Sun, Vincent Vargas, Menglu Wang, Samuel Watson, Wendelin Werner, David Wilson, and Hao Wu.

We would also like to thank the Isaac Newton Institute (INI) for Mathematical Sciences, Cambridge, for its support and hospitality during the program on Random Geometry where part of this work was completed. J.M.'s work was also partially supported by DMS-1204894 and S.S.'s work was also partially supported by DMS-1209044, a fellowship from the Simons Foundation, and EPSRC grants EP/L018896/1 and EP/I03372X/1.

1 Introduction

1.1 Overview

The Liouville quantum gravity (LQG) sphere [DS11, She15, DMS14, MS15b, DKRV14, AHS15] and the Brownian map (TBM) [MM06, LGP08, LG13, Mie13, LG14, MS15b] can both be understood as measures on the space of measure-endowed, sphere-homeomorphic surfaces. The definition of LQG involves a real parameter γ, and it has long been believed that when $\gamma = \sqrt{8/3}$, the LQG sphere should be equivalent to TBM. However, the LQG sphere comes equipped with a conformal structure and TBM comes equipped with a metric space structure, and it is far from obvious how to endow either of these objects with the *other's* structure.

This paper is the first in a three part series (also including [MS16a, MS16b]) that will provide a robust unification of $\sqrt{8/3}$-LQG and TBM. Over the course of these three papers, we will show the following:

1. An instance of the $\sqrt{8/3}$-LQG sphere a.s. comes with a **canonical metric space structure**, and the resulting measure-endowed metric space agrees in law with TBM.

2. Given an instance of TBM, the $\sqrt{8/3}$-LQG sphere that generates it is a.s. uniquely determined. This implies that an instance of TBM a.s. comes with a canonical (up to Möbius transformation) embedding in the Euclidean sphere. In other words:

3. An instance of TBM a.s. comes with a **canonical conformal structure** and the resulting measure-endowed conformal sphere agrees in law with the $\sqrt{8/3}$-LQG sphere.

4. The canonical (up to Möbius transformation) embedding of TBM (with its intrinsic metric) into the Euclidean sphere (with the Euclidean metric) is a.s. Hölder continuous with Hölder continuous inverse. This in particular implies that a.s. all geodesics in TBM are sent to Hölder continuous curves in the Euclidean sphere.

In [MS16a, MS16b], we will also extend these results to infinite volume surfaces (the so-called Brownian plane [CL12] and the $\sqrt{8/3}$-LQG quantum cone [She15, DMS14]) and to surfaces with boundary (the Brownian disk and its LQG analog).

Before we begin to explain how these results will be proved, let us describe one reason one might expect them to be true. Both TBM and the $\sqrt{8/3}$-LQG sphere are known to be $n \to \infty$ scaling limits of the uniformly random planar map with n edges, albeit w.r.t. different topologies.[1] One should therefore be able to use a compactness argument to show that the uniformly random planar map has a scaling limit (at least subsequentially) in the product of these topologies, and that this scaling limit is a *coupling* of TBM and the $\sqrt{8/3}$-LQG sphere. It is not obvious that, in this coupling, the instance of TBM and the instance of the $\sqrt{8/3}$-LQG sphere a.s. uniquely determine one another. But it seems reasonable to guess that this would be the case. And it is at least conceivable that one could prove this through a sophisticated analysis of the planar maps themselves (e.g., by showing that pairs of random planar maps are highly likely to be close in one topology if and only if they are close in the other topology).

Another reason to guess that an LQG sphere should have a canonical metric structure, and that TBM should have a canonical conformal structure, is that it is rather easy to formulate reasonable sounding *conjectures* about how a metric on an LQG sphere might be obtained as limit of *approximate* metrics, or how a conformal structure on TBM might be obtained as a limit of *approximate* conformal structures. For example, the peanosphere construction of [DMS14] gives a space-filling curve on the LQG sphere; one might divide space into regions traversed by length-δ increments of time, declare two such regions adjacent if they intersect, and conjecture that the corresponding graph distance (suitably rescaled) converges to a continuum distance as $\delta \to 0$. Similarly, an instance of TBM comes with a natural space-filling curve; one can use this to define a graph structure as above, embed the graph in the Euclidean sphere using circle packing (or some other method thought to respect conformal structure), and conjecture that as $\delta \to 0$ these embeddings converge to a canonical (up to Möbius transformation) embedding of TBM in the Euclidean sphere. In both of these cases, the approximating graph can be constructed in a simple way (in terms of Brownian motion or the Brownian snake) and could in principle be studied directly.

[1] TBM is the scaling limit w.r.t. the Gromov-Hausdorff topology on metric spaces [LGP08, LG13, Mie13, LG14]. The $\sqrt{8/3}$-LQG sphere (decorated by CLE_6) is the scaling limit of the uniformly random planar map (decorated by critical percolation) w.r.t. the so-called peanosphere topology, as well as a stronger topology that encodes loop lengths and intersection patterns (see [She11, DMS14], the forthcoming works [GS15a, GS15b, GM16], and the brief outline in [MS15a]).

The current series of papers will approach the problem from a completely different direction, which we believe to be easier and arguably more enlightening than the approaches suggested above. Instead of using approximations of the sort described above, we will use a combination of the quantum Loewner evolution (QLE) ideas introduced in [MS13b], TBM analysis that appears in [MS15a], and the $\sqrt{8/3}$-LQG sphere analysis that appears in [MS15b]. There are approximations involved in defining the relevant form of QLE, but they seem to respect the natural symmetries of the problem in a way that the approximation schemes discussed above do not. In particular, our approach will allow us to take full advantage of an exact relationship between the LQG disks "cut out" by SLE_6 and those cut out by a metric exploration.

In order to explain our approach, let us introduce some notation. If (S, d) is a metric space, like TBM, and $x \in S$ then we let $B(x, r)$ denote the radius r ball centered at x. If the space is homeomorphic to $\mathbf{S}^2$ and comes with a distinguished "target" point y, then we let $B^\bullet(x, r)$ denote the *filled* metric ball of radius r, i.e., the set of all points that are disconnected from y by $\overline{B(x, r)}$. Note that if $0 \leq r < d(x, y)$, then the complement of $B^\bullet(x, r)$ contains y and is homeomorphic to the unit disk $\mathbf{D}$.

The starting point of our approach is to let x and y be points on an LQG-sphere and to define a certain "growth process" growing from x to y. We assume that x and y are "quantum typical," i.e., that given the LQG-sphere itself, the points x and y are independent samples from the LQG measure on that sphere. The growth process is an increasing family of closed subsets of the LQG-sphere, indexed by a time parameter t, which we denote by $\Gamma_t = \Gamma_t^{x \to y}$. Ultimately, the set Γ_t will represent the filled metric ball $B^\bullet(x, t)$ corresponding to an appropriately defined metric on the LQG-sphere (when y is taken to be the distinguished target point). However, we will get to this correspondence somewhat indirectly. Namely, we will *first* define $\Gamma_t = \Gamma_t^{x \to y}$ as a random growth process (for quantum typical points x and y) and only show *a posteriori* that there is a metric for which the Γ_t thus defined are a.s. the filled metric balls.

As presented in [MS13b], the idea behind this growth process (whose discrete analog we briefly review and motivate in Section 3) is that one should be able to "reshuffle" the SLE_6 decorated quantum sphere in a particular way in order to obtain a growth process on a LQG surface that hits points in order of their distance from the origin. This process is a variant of the $\text{QLE}(8/3, 0)$ process originally constructed in [MS13b] by starting with an SLE_6 process and then "resampling" the tip location at small capacity time increments to obtain a type of first passage percolation on top of a $\sqrt{8/3}$-LQG surface. The form of $\text{QLE}(8/3, 0)$ that we use here differs from that given in [MS13b] in that we will resample the tip at "quantum natural time" increments as defined in [DMS14] (i.e., time steps which are intrinsic to the surface rather than to its specific choice of embedding). We expect that these two constructions are in fact equivalent, but we will not establish that fact here.

As discussed in [MS13b], the growth process $\text{QLE}(8/3, 0)$ can in some sense be understood as a continuum analog of the Eden model. The idea explained there is that in some

small-increment limiting sense, the (random) Eden model growth should correspond to (deterministic) metric growth. In fact, a version of this statement for random planar maps has recently been verified in [CL14a], which shows that on a random planar map, the random metric associated with an Eden model (or first passage percolation) instance closely approximates graph distance.

Once we have defined the growth process for quantum typical points x and y, we will define the quantity $d_{\mathcal{Q}}(x, y)$ to be the amount of time it takes for a QLE growth process to evolve from x to y. This $d_{\mathcal{Q}}$ is a good candidate to be a distance function, at least for those x and y for which it is defined. However, while our initial construction of QLE will produce the joint law of the doubly marked LQG surface and the growth process Γ, it will not be obvious from this construction that $d_{\mathcal{Q}}(x, y) = d_{\mathcal{Q}}(y, x)$ almost surely. In fact, it will not even be obvious that the growth process is a.s. *determined* by the LQG sphere and the (x, y) pair, so we will *a priori* have to treat $d_{\mathcal{Q}}(x, y)$ as a random variable whose value might depend on more than the LQG-surface and the (x, y) pair.

The bulk of the current paper is dedicated to showing that if one first samples a $\sqrt{8/3}$-LQG sphere, and then samples $x_1, x_2, \ldots$ as i.i.d. samples from its LQG measure, and then samples conditionally independent growth processes from each x_i to each x_j, then it is almost surely the case that the $d_{\mathcal{Q}}$ defined from these growth processes is a metric, and that this metric is determined by the LQG sphere and the points, as stated in Theorem 1.1 below.

Both the $\sqrt{8/3}$-LQG sphere and TBM have some natural variants that differ in how one handles the issues of total area and special marked points; these variants are explained for example in [MS15a, MS15b]. On both sides, there is a natural unit area sphere measure $d\mathcal{S}$ in which the total area measure is a.s. one. On both sides, one can represent a sphere of arbitrary positive and finite area by a pair $(\mathcal{S}, A)$, where $\mathcal{S}$ is a unit area sphere and A is a positive real number. The pair represents the unit area sphere $\mathcal{S}$ except with area scaled by a factor of A and distance (where defined) scaled by a factor of $A^{1/4}$. On both sides it turns out to be natural to define infinite measures on quantum spheres such that the "total area" marginal has the form $A^{\beta} dA$ for some β. In particular, on both sides, one can define a natural "grand canonical" quantum sphere measure on spheres with k marked points (see the notation in Section 1.3). Sampling from this infinite measure amounts to

1. first sampling a unit area sphere $\mathcal{S}$,
2. then sampling k marked points i.i.d. from the measure on $\mathcal{S}$,
3. then independently selecting A from the infinite measure $A^{k-7/2} dA$ and rescaling the sphere's area by a factor of A (and distance by a factor of $A^{1/4}$).

Theorem 1.1, stated below, applies to all of these variants. Recall that in the context of an infinite measure, *almost surely* means outside a set of measure zero.

Theorem 1.1. *Suppose that $\mathcal{S}$ is an instance of the $\sqrt{8/3}$-quantum sphere, as defined in [DMS14, MS15b] (either the unit area version or the "grand canonical" version involving one or more distinguished points). Let (x_n) be a sequence of points chosen independently from the quantum area measure on $\mathcal{S}$. Then it is a.s. the case that for each pair x_i, x_j, the quantity $d_{\mathcal{Q}}(x_i, x_j)$ is uniquely determined by $\mathcal{S}$ and the points x_i and x_j. Moreover, it is a.s. the case that*

1. *$d_{\mathcal{Q}}(x_i, x_j) = d_{\mathcal{Q}}(x_j, x_i) \in (0, \infty)$ for all distinct i and j, and*
2. *$d_{\mathcal{Q}}$ satisfies the (strict) triangle inequality $d_{\mathcal{Q}}(x_i, x_k) < d_{\mathcal{Q}}(x_i, x_j) + d_{\mathcal{Q}}(x_j, x_k)$ for all distinct i, j, and k.*

The fact that the triangle inequality in Theorem 1.1 is a.s. strict implies that *if* the metric $d_{\mathcal{Q}}$ can be extended to a geodesic metric on the entire LQG-sphere (something we will establish in the subsequent paper [MS16a]) then in this metric it is a.s. the case that none of the points on the countable sequence lies on a geodesic between two other points in the sequence. This is unsurprising given that, in TBM, the measure of a geodesic between two randomly chosen points is almost surely zero. (This is well known and essentially immediate from the definition of TBM; see [MS15a] for some discussion of this point.)

The construction of the metric in Theorem 1.1 is "local" in the sense that it only requires that the field near any given point is absolutely continuous with respect to a $\sqrt{8/3}$-LQG sphere. In particular, Theorem 1.1 yields a construction of the metric on a countable, dense subset of any $\sqrt{8/3}$-LQG surface chosen i.i.d. from the quantum measure. Moreover, the results of the later papers [MS16a, MS16b] also apply in this generality, which allows one to define geodesic metrics on other $\sqrt{8/3}$-LQG surfaces, such as the torus constructed in [DRV15].

The proof of Theorem 1.1 is inspired by a closely related argument used in a paper by the second author, Sam Watson, and Hao Wu (still in preparation) to define a metric on the set of loops in a CLE_4 process. To briefly sketch how the proof goes, suppose that we choose a $\sqrt{8/3}$-LQG sphere $\mathcal{S}$ with marked points x and y, and then choose a growth process Γ from x to y and a conditionally independent growth process $\overline{\Gamma}$ from y to x. We also let U be chosen uniformly in $[0,1]$ independently of everything else. Let Θ be the joint distribution of $(\mathcal{S}, x, y, \Gamma, \overline{\Gamma}, U)$. Since the natural measure on $\sqrt{8/3}$-LQG spheres is an infinite measure, so is Θ. However, we can make sense of Θ conditioned on $\mathcal{S}$ as a probability measure. Given $\mathcal{S}$, we have that $d_{\mathcal{Q}}(x, y)$ and $d_{\mathcal{Q}}(y, x)$ are well defined as random variables denoting the respective time durations of Γ and $\overline{\Gamma}$. As discussed above, we interpret $d_{\mathcal{Q}}(x, y)$ (resp. $d_{\mathcal{Q}}(y, x)$) as a measure of the distance from x to y (resp. y to x). Write $\Theta^{x \to y}$ for the weighted measure $d_{\mathcal{Q}}(x, y) d\Theta$. In light of the uniqueness of Radon-Nikodym derivatives, in order to show that $d_{\mathcal{Q}}(x, y) = d_{\mathcal{Q}}(y, x)$ almost surely, it will suffice to show that $\Theta^{x \to y} = \Theta^{y \to x}$.



Figure 1.1: A "figure-8" consisting of meeting QLE$(8/3, 0)$ processes which illustrates the statement of Lemma 1.2, the main input into the proof of Theorem 1.1. The blue (resp. red) arcs contained in the illustration of $\Gamma|_{[0,\tau]}$ (resp. $\overline{\Gamma}|_{[0,\overline{\tau}]}$) represent the regions which have been cut off from y (resp. x).

The main input into the proof of this is Lemma 1.2, stated below (which will later be restated slightly more precisely and proved as Theorem 7.1). Suppose we sample $(\mathcal{S}, x, y, \Gamma, \overline{\Gamma}, U)$ from $\Theta^{x\to y}$, then let $\tau = U d_{\mathcal{Q}}(x, y)$ — so that τ is uniform in $[0, d_{\mathcal{Q}}(x, y)]$, and then define $\overline{\tau} = \inf\{t \geq 0 : \overline{\Gamma}_t \cap \Gamma_\tau \neq \emptyset\}$. Both $\Gamma|_{[0,\tau]}$ and $\overline{\Gamma}|_{[0,\overline{\tau}]}$ are understood as growth processes *truncated* at random times, as illustrated in Figure 1.1. We also let $\overline{\sigma} = U d_{\mathcal{Q}}(y, x)$ and $\sigma = \inf\{t \geq 0 : \Gamma_t \cap \overline{\Gamma}_{\overline{\sigma}} \neq \emptyset\}$. Under $\Theta^{y\to x}$, we have that $\overline{\sigma}$ is uniform in $[0, d_{\mathcal{Q}}(y, x)]$.

Lemma 1.2. *Using the definitions above, the $\Theta^{x\to y}$-induced law of $(\mathcal{S}, x, y, \Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]})$ is equal to the $\Theta^{y\to x}$-induced law of $(\mathcal{S}, x, y, \Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]})$.*

The proof of Lemma 1.2 is in some sense the heart of the paper. It is established in Section 7, using tools developed over several previous sections, which in turn rely on the detailed understanding of SLE_6 processes on $\sqrt{8/3}$-LQG spheres developed in [DMS14, MS15b] as well as in [MS12a, MS12b, MS12c, MS13a]. We note that the intuition behind this symmetry was also sketched at the end of [MS15a] in the context of TBM.

To derive $\Theta^{x\to y} = \Theta^{y\to x}$ from Lemma 1.2, it will suffice to show the following:

Lemma 1.3. *The $\Theta^{x\to y}$ conditional law of Γ, $\overline{\Gamma}$ given $(\mathcal{S}, x, y, \Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]})$ is the same as the $\Theta^{y\to x}$ conditional law of Γ, $\overline{\Gamma}$ given $(\mathcal{S}, x, y, \Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]})$.*

Intuitively, sampling from either conditional law should amount to just continuing the evolution of Γ and $\overline{\Gamma}$ on $\mathcal{S}$, beyond their given stopping times, independently of each other. However, it will take some work to make this intuition precise. The remainder of this subsection provides a short overview of the argument.

Because of the way that $\Theta^{x\to y}$ was constructed, one can show quite easily that the $\Theta^{x\to y}$ conditional law of Γ *given* the five-tuple $(\mathcal{S}, x, y, \Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]})$ is a function of the four-tuple $(\mathcal{S}, x, y, \Gamma|_{[0,\tau]})$. (The existence of regular conditional probabilities follows from the fact that the Gaussian free field and associated growth processes can be defined as random variables in a standard Borel space; see further discussion for example in [SS13, MS13b].) The proof of Lemma 1.3 will be essentially done once we establish the following claim: the $\Theta^{y\to x}$ conditional law of Γ *given* the five-tuple $(\mathcal{S}, x, y, \Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]})$ is described by the same function applied to the four-tuple $(\mathcal{S}, x, y, \Gamma|_{[0,\sigma]})$. Indeed, a symmetric argument implies an analogous statement about the conditional law of $\overline{\Gamma}$ under $\Theta^{x\to y}$ and $\Theta^{y\to x}$ given the corresponding five-tuple. Lemma 1.3 will follow readily from this symmetric pair of statements and the *a priori* conditional independence of Γ and $\overline{\Gamma}$ given $(\mathcal{S}, x, y)$.

The claim stated just above may appear to be obvious, but there is still some subtlety arising from the fact that τ and σ are not defined in symmetric ways *a priori*, and in fact σ is a complicated stopping time for Γ (which depends both on $(\mathcal{S}, x, y)$ and on the additional randomness encoded in $\overline{\Gamma}$ and $\overline{\sigma}$), and we have not proved anything like a strong Markov property that would hold for arbitrary stopping times. To deal with this rather technical point, we will approximate both τ and σ by stopping times that a.s. take only countably many values, noting that the required Markov property is easy to derive for such stopping times. To prove that the conditional laws computed using these approximations converge to the appropriate limits, we will check that for any bounded function F the process $\mathbf{E}_\Theta\big[F(\Gamma)\,|\,(\mathcal{S}, x, y, \Gamma|_{[0,r]})\big]$ is a.s. continuous (as a function of r) at the times τ and σ, respectively. Since these processes are martingales, they a.s. have only countably many discontinuities; it will thus suffice to prove that when $(\mathcal{S}, x, y, \Gamma, \overline{\Gamma})$ is given and τ (resp. $\overline{\sigma}$) is chosen uniformly from $[0, d_{\mathcal{Q}}(x, y)]$ (resp. $[0, d_{\mathcal{Q}}(y, x)]$) the probability that either τ or σ assumes any *fixed* value is zero. This is obviously true for τ, and it follows for σ because the map $\overline{\sigma} \to \sigma$ (a random function that depends on $(\mathcal{S}, x, y, \Gamma, \overline{\Gamma})$) is non-increasing by definition, and the symmetry property implies that Γ a.s. first hits $\overline{\Gamma}_{\overline{\sigma}}$ at exactly time σ, which implies that there a.s. cannot be a positive interval of $\overline{\sigma}$ values on which σ is constant.

We will see *a posteriori* that $\tau = d_{\mathcal{Q}}(x, y) - \overline{\tau}$, which we will prove by using the fact that $d_{\mathcal{Q}}(x, y) = d_{\mathcal{Q}}(y, x)$ and the symmetry of Lemma 1.2, which implies that both $\overline{\tau}$ and τ are $\Theta^{x\to y}$-conditionally uniform on $[0, d_{\mathcal{Q}}(x, y)]$, once $(\mathcal{S}, x, y)$ is given. We will also use this fact to derive the triangle inequality. Note that if z is a third point and we are working on the event that $d_{\mathcal{Q}}(x, z) < d_{\mathcal{Q}}(x, y)$, then $\Gamma_{d_{\mathcal{Q}}(x,z)}$ and $\overline{\Gamma}_{d_{\mathcal{Q}}(y,z)}$ must intersect each other, at least at the point z. In fact, it will not be hard to see that a.s. for some $\epsilon > 0$ the processes $\Gamma_{d_{\mathcal{Q}}(x,z)}$ and $\overline{\Gamma}_{d_{\mathcal{Q}}(y,z)-\epsilon}$ still intersect. This implies that if $\tau = d_{\mathcal{Q}}(x, z)$ then $\overline{\tau} \le d_{\mathcal{Q}}(y, z) - \epsilon < d_{\mathcal{Q}}(y, z)$. Plugging in $\overline{\tau} = d_{\mathcal{Q}}(x, y) - \tau = d_{\mathcal{Q}}(x, y) - d_{\mathcal{Q}}(x, z)$, we obtain the strict triangle inequality.

1.2 Further observations and sequel overview

In the course of establishing Theorem 1.1, it will also become clear (almost immediately from the definitions) that the growth process from x_i to x_j and the growth process from x_i to x_k almost surely agree up until some random time at which x_j and x_k are first separated from each other, after which the two processes evolve independently. Thus one can describe the full collection of growth processes from x_i to all the other points in terms of a single "branching" growth process with countably many "branch times" (i.e., times at which some x_j and x_k are separated for the first time).

It will also become clear from our construction that when exploring from a marked point x to a marked point y, one can make sense of the length of $\partial B^\bullet(x,t)$, and that as a process indexed by t this evolves as the time-reversal of an excursion of a continuous state branching process (CSBP), with the jumps in this process corresponding to branch times.[2] We will review the definition of a CSBP in Section 2.1. Letting y vary over all of the points x_j, one obtains a branching version of a time-reversed CSBP excursion, and it will become clear that the law of this branching process agrees with the analogous law for TBM, as explained in [MS15a].

All of this suggests that we are well on our way to establishing the equivalence of the $\sqrt{8/3}$-LQG sphere and TBM. As further evidence in this direction, note that it was established in [MS15a] that the time-reversed branching process (together with a countable set of real numbers indicating where along the boundary each "pinch point" occurs) contains all of the information necessary to reconstruct an instance of the entire Brownian map. That is, given a complete understanding of the exploration process rooted at a single point, one can a.s. reconstruct the distances between all pairs of points. This suggests (though we will not make this precise until the subsequent paper [MS16a]) that, given the information described in the QLE branching process provided in this paper, one should also be able to recover an entire Brownian map instance.

In order to finish the project, the program in [MS16a] will be to

1. Derive some continuity estimates and use them to show that $d_{\mathcal{Q}}$ a.s. extends uniquely to a metric defined on the entire LQG sphere, and to establish Hölder continuity for the identify map from the sphere endowed with the Euclidean metric to the sphere endowed with the random metric, and then

2. Learn enough about the geodesics within the so-called metric net (as defined in [MS15b]) to allow us to show that the random metric satisfies the properties that are shown in [MS15b] to characterize TBM.

[2] We will see in [MS16a] that if one defines a quantum-time QLE$(8/3, 0)$ on an infinite volume LQG surface, namely a $\sqrt{8/3}$ quantum cone, then the evolution of the boundary length is a process that matches the one described by Krikun (discrete) [Kri05] and Curien and Le Gall [CL14b] for the Brownian plane [CL12].

This will imply that the metric space described by $d_{\mathcal{Q}}$ has the law of TBM and, moreover, that the instance of TBM is a.s. determined by the underlying $\sqrt{8/3}$-LQG sphere. The program in [MS16b] will be to prove that in the coupling between TBM and the $\sqrt{8/3}$-LQG sphere, the former a.s. determines the latter, i.e., to show that an instance of TBM a.s. has a canonical embedding into the sphere. Thus we will have that the $\sqrt{8/3}$-LQG sphere and TBM are equivalent in the sense that an instance of one a.s. determines the other. The ideas used in [MS16b] will be related to the arguments used in [DMS14] to show that an instance of the peanosphere a.s. has a canonical embedding.

1.3 Prequel overview

As noted in Section 1.1, both TBM and the $\sqrt{8/3}$-LQG sphere have natural infinite volume variants that in some sense correspond to grand canonical ensembles decorated by some fixed number of marked points. In this paper, because we deal frequently with exploration processes from one marked point to another, we will be particularly interested in the natural infinite measures on *doubly* marked spheres. We recall that

1. In [MS15a] this natural measure on doubly marked Brownian map spheres with two marked points is denoted μ^2_{SPH} (and more generally μ^k_{SPH} refers to the measure with k marked points).
2. In [MS15b] the natural measure on doubly marked $\sqrt{8/3}$-LQG spheres is denoted by $\mathsf{M}^2_{\text{SPH}}$.

As noted in Section 1.1, in both cases, the law of the overall area is given (up to a multiplicative constant) by $A^{-3/2}dA$. In both cases, the conditional law of the surface *given* A is that of a sample from a probability measure on unit area surfaces (with the measure rescaled by a factor of A, and distance rescaled by $A^{1/4}$ — though of course distance is not *a priori* defined on the LQG side). We remark that in much of the literature on TBM the unit area measure is the primary focus of attention (and it is denoted by $\mu^{A=1}_{\text{SPH}}$ in [MS15a]).

The paper [MS15b] explains how to explore a doubly marked surface $(\mathcal{S}, x, y)$ sampled from $\mathsf{M}^2_{\text{SPH}}$ with an SLE_6 curve drawn from x to y. The paper [MS15a] explains how to explore a doubly marked surface $(\mathcal{S}, x, y)$ sampled from μ^2_{SPH} by exploring the so-called "metric net," which consists of the set of points that lie on the outer boundary of $B^\bullet(x, r)$ for some $r \in [0, d(x, y)]$. (We are abusing notation slightly here in that $\mathcal{S}$ represents a quantum surface in the first case and a metric space in the second, and these are *a priori* different types of objects.) In both cases, the exploration/growth procedure "cuts out" a countable collection of disks, each of which comes with a well defined boundary length. Also in both cases, the process that encodes the boundary length corresponds (up to time change) to the set of jumps in the *time-reversal* of a 3/2-stable Lévy excursion with only positive jumps. Moreover, in both cases, the boundary length of each disk "cut

out" is encoded by the length of the corresponding jump in the time-reversed 3/2-stable Lévy excursion. Finally, in both cases, the disks can be understood as conditionally independent measure-endowed random surfaces, given their boundary lengths.

The intuitive reason for the similarities between these two types of explorations is explained in the QLE paper [MS13b], and briefly reviewed in Section 3. The basic idea is that in the discrete models involving triangulations, the conditional law of the unexplored region (the component containing y) does not depend on the rule one uses to decide which triangle to explore next; if one is exploring via the Eden model, one picks a random location on the boundary to explore, and if one is exploring a percolation interface, one explores along a given path. The law of the set of disks cut out by the exploration is the same in both cases.

The law of a "cut out" disk, given that its boundary length is L, is referred to as μ^L_{DISK} in [MS15a]. If one explores up to some stopping time before encountering y, then the conditional law of the unexplored region containing y is that of a *marked* Brownian disk with boundary length L (here y is the marked point), and is referred to as $\mu^{1,L}_{\text{DISK}}$ in [MS15a]. It is not hard to describe how these two measures are related. If one forgets the marked point y, then both μ^L_{DISK} and $\mu^{1,L}_{\text{DISK}}$ describe probability measures on the space of quantum disks; and from this perspective, the Radon-Nikodym derivative of $\mu^{1,L}_{\text{DISK}}$ w.r.t. μ^L_{DISK} is given (up to multiplicative constant) by the total surface area. Given the quantum disk sampled from $\mu^{1,L}_{\text{DISK}}$, the conditional law of the marked point y is that of a sample from the quantum measure on the surface.

Precisely analogous statements are given in [MS15b] for the SLE_6 exploration of a sample from $\mathsf{M}^2_{\text{SPH}}$.[3] The following objects are shown in [MS15b] well defined, and are analogous to objects produced by the measures μ^L_{DISK} and $\mu^{1,L}_{\text{DISK}}$ in [MS15a]:

1. **The $\sqrt{8/3}$-LQG disk with boundary length L.** This is a random quantum surface whose law is the conditional law of a surface cut out by the SLE_6 exploration, given only its boundary length (and not its embedding in the larger surface).

2. **The *marked* $\sqrt{8/3}$-LQG disk with boundary length L.** This is a random quantum surface whose law is obtained by weighting the *unmarked* law by total area, and letting the conditional law of y given the surface be that of a uniformly random sample from the area measure (normalized to be a probability measure). It represents the conditional law of the unexplored quantum component containing y.

Proposition 1.4. *Consider a doubly marked $\sqrt{8/3}$-LQG sphere decorated by an independent whole plane* SLE_6 *path η' from its first marked point x to its second marked*

[3] These results are in turn consequences of the fact, derived by the authors and Duplantier in an infinite volume setting in [DMS14], that one can weld together two so-called Lévy trees of $\sqrt{8/3}$-LQG disks to produce a new $\sqrt{8/3}$-LQG surface decorated by an independent SLE_6 curve that represents the interface between the two trees.



Figure 1.2: **Left:** A whole-plane SLE_6 process η' from x to y is drawn on top of a doubly-marked $\sqrt{8/3}$-LQG sphere $(\mathcal{S}, x, y)$. The grey region represents the part of $\mathcal{S}$ that $\eta'([0, k\delta])$ has disconnected from y at time $k\delta$, and $T_{k\delta}$ is the outer boundary of that region. We note that $T_{k\delta}$ has cut points because whole-plane SLE_6 has cut points. **Middle:** Green region indicates the additional area cut off from y by $\eta'([0, (k+1)\delta])$, and $T_{(k+1)\delta}$ is the outer boundary $\eta'([0, (k+1)]\delta])$. **Right:** It is conceptually useful to imagine that we "cut" along both $T_{k\delta}$ and $T_{(k+1)\delta}$ to produce a beaded quantum surface $\mathcal{N}_k$ (a so-called "necklace") attached to a loose "string" whose length is the length of $T_{(k+1)\delta} \cap T_{k\delta}$. The inner circle is understood to have total length equal to the length of $T_{k\delta}$, and the outer boundary has total length equal to the length of $T_{(k+1)\delta}$.

point y. We consider η' to be parameterized by its quantum natural time. Fix an $s > 0$ and let T_s denote the outer boundary of the closed set $\eta'([0, s])$, i.e., the boundary of the y-containing component of the complement of $\eta'([0, s])$. Then the conditional law of the y-containing region (given that its boundary length is L_s) is that of a marked $\sqrt{8/3}$-LQG disk with boundary length L_s. In particular, since this law is rotationally invariant, the overall law of the surface is unchanged by the following random operation: "cut" along T_s, rotate the disk cut out by a uniformly random number in $[0, L_s]$, and then weld this disk back to the beaded quantum surface $\eta'([0, s])$ (again matching up quantum boundary lengths).

It is natural to allow s to range over integer multiples of a constant δ. As illustrated in Figure 1.2, we let $\mathcal{N}_k$ denote the "necklace" described by the union $T_{k\delta} \cup T_{(k+1)\delta} \cup \eta'\Big([k\delta, (k+1)\delta]\Big)$, which we interpret as a *beaded quantum surface* (see [DMS14]) attached to a "string" of some well defined length. Applying the above resampling for each integer multiple of δ corresponds to "reshuffling" these necklaces in the manner depicted in Figure 3.4.

Proposition 1.5. *Fix $\delta > 0$ and apply the random rotation described in Proposition 1.4 for each s that is an integer multiple of δ. Taking any subsequential limit as $\delta \to 0$, we*

obtain a coupling of a $\sqrt{8/3}$-quantum sphere with a growth process on that sphere, such that the law of the ordered set of disks cut out by that process is the same as in the SLE_6 *case.*

The growth process obtained this way is what we will call the *quantum natural time* $\mathrm{QLE}(8/3, 0)$ (as opposed to the *capacity time* $\mathrm{QLE}(8/3, 0)$ process described in [MS13b], which we expect but do not prove to be equivalent to the quantum time version). As already noted in Section 1.1, we will make extensive use of quantum natural time $\mathrm{QLE}(8/3, 0)$ in this paper. When we use the term $\mathrm{QLE}(8/3, 0)$ without a qualifier, we will mean the quantum natural time variant.

Let us highlight one subtle point about this paper. Although we *a priori* construct only *subsequential limits* for the growth process $\mathrm{QLE}(8/3, 0)$ using the procedure described in Proposition 1.5, we ultimately show that the metric $d_{\mathcal{Q}}$ defined on a countable sequence of i.i.d. points (x_n) chosen from the quantum measure does not depend on the particular choice of subsequence. Once we know this metric we know, for each $t \geq 0$ and each x and y in (x_n), which points from the set (x_n) lie in the set $\Gamma_t^{x\to y}$. Since $\Gamma_t^{x\to y}$ is closed, we would expect it to be given by precisely the closure of this set of points, which would imply that the growth process described in Proposition 1.5 is a.s. defined as a true (non-subsequential) limit. This would follow immediately if we knew, say, that $\Gamma_t^{x\to y}$ was a.s. the closure of its interior. However, we will not prove in this paper that this is the case. That is, we will not rule out the possibility that the boundary of $\Gamma_t^{x\to y}$ contains extra "tentacles" that possess zero quantum area and somehow fail to intersect any of the (x_n) values. Ruling out this type of behavior will be part of the program in [MS16a], where we establish a number of continuity estimates for $\mathrm{QLE}(8/3, 0)$ and $d_{\mathcal{Q}}$. Upon showing this, we will be able to remove the word "subsequential" from the statement of Proposition 1.5.

1.4 Outline

The remainder of this article is structured as follows. In Section 2 we review preliminary facts about continuous state branching processes, quantum surfaces, and conformal removability. In Section 3, we recall some of the discrete constructions on random planar triangulations that appeared in [MS13b], which we use to explain and motivate our continuum growth processes. In particular, we will recall that on these triangulated surfaces *random metric explorations* are in some sense "reshufflings" of *percolation explorations*, and in Section 4 we construct quantum-time $\mathrm{QLE}(8/3, 0)$ using an analogous reshuffling of SLE_6. In Section 5 we establish a certain symmetry property for continuum percolation explorations (SLE_6) on $\sqrt{8/3}$-LQG surfaces (a precursor to the main symmetry result we require). Then in Section 6 we will give the construction of the quantum natural time variant of $\mathrm{QLE}(8/3, 0)$. In Section 7, we establish the main symmetry result we require, and in Section 8 we will finish the proof of Theorem 1.1.

2 Preliminaries

2.1 Continuous state branching processes

We will now review some of the basic properties of continuous state branching processes (CSBPs) and their relationship to Lévy processes. CSBPs will arise in this article because they describe the time-evolution of the quantum boundary length of the boundary of a QLE$(8/3, 0)$. We refer the reader to [Ber96] for an introduction to Lévy processes and to [LG99, Kyp06] for an introduction to CSBPs.

A CSBP with branching mechanism ψ (or ψ-CSBP for short) is a Markov process Y on $\mathbf{R}_+$ whose transition kernels are characterized by the property that

$$\mathbf{E}[\exp(-\lambda Y_t) \,|\, Y_s] = \exp(-Y_s u_{t-s}(\lambda)) \quad \text{for all} \quad t > s \geq 0 \tag{2.1}$$

where $u_t(\lambda)$, $t \geq 0$, is the non-negative solution to the differential equation

$$\frac{\partial u_t}{\partial t}(\lambda) = -\psi(u_t(\lambda)) \quad \text{for} \quad u_0(\lambda) = \lambda. \tag{2.2}$$

Let

$$\Phi(q) = \sup\{\theta \geq 0 : \psi(\theta) = q\} \tag{2.3}$$

and let

$$\zeta = \inf\{t \geq 0 : Y_t = 0\} \tag{2.4}$$

be the extinction time for Y. Then we have that [Kyp06, Corollary 10.9]

$$\mathbf{E}\left[e^{-q\int_0^\zeta Y_s ds}\right] = e^{-\Phi(q)Y_0}. \tag{2.5}$$

A ψ-CSBP can be constructed from a Lévy process with only positive jumps and vice-versa [Lam67] (see also [Kyp06, Theorem 10.2]). Namely, suppose that X is a Lévy process with Laplace exponent ψ. That is, if $X_0 = x$ then we have that

$$\mathbf{E}[e^{-\lambda(X_t - x)}] = e^{\psi(\lambda)t}.$$

Let

$$s(t) = \int_0^t \frac{1}{X_u} du \quad \text{and} \quad s^*(t) = \inf\{r > 0 : s(r) > t\}. \tag{2.6}$$

Then the time-changed process $Y_t = X_{s^*(t)}$ is a ψ-CSBP. Conversely, if Y is a ψ-CSBP and we let

$$t(s) = \int_0^s Y_u du \quad \text{and} \quad t^*(s) = \inf\{r > 0 : t(r) > s\} \tag{2.7}$$

then $X_s = Y_{t^*(s)}$ is a Lévy process with Laplace exponent ψ.

We will be interested in the particular case that $\psi(u) = u^\alpha$ for $\alpha \in (1, 2)$. For this choice, we note that

$$u_t(\lambda) = \left(\lambda^{1-\alpha} + (\alpha - 1)t\right)^{1/(1-\alpha)}. \tag{2.8}$$

2.2 Quantum surfaces

Suppose that h is an instance of (a form of) the Gaussian free field (GFF) on a planar domain D and $\gamma \in [0,2)$ is fixed. Then the γ-LQG surface associated with h is described by the measure μ_h which is formally given by $e^{\gamma h(z)}dz$ where dz denotes Lebesgue measure on D. Since the GFF h does not take values at points (it is a random variable which takes values in the space of distributions), it takes some care to make this definition precise. One way of doing so is to let, for each $\epsilon > 0$ and $z \in D$ such that $B(z,\epsilon) \subseteq D$, $h_\epsilon(z)$ be the average of h on $\partial B(z,\epsilon)$ (see [DS11, Section 3] for more on the circle average process). The process $(z,\epsilon) \mapsto h_\epsilon(z)$ is jointly continuous in (z,ϵ) and one can define $e^{\gamma h(z)}dz$ to be the weak limit as $\epsilon \to 0$ along negative powers of 2 of $\epsilon^{\gamma^2/2}e^{\gamma h_\epsilon(z)}dz$ [DS11]; the normalization factor $\epsilon^{\gamma^2/2}$ is necessary for the limit to be non-trivial. We will often write μ_h for the measure $e^{\gamma h(z)}dz$. In the case that h has free boundary conditions, one can also construct the natural boundary length measure $\nu_h = e^{\gamma h(z)/2}dz$ in a similar manner.

The regularization procedure used to construct μ_h leads to the following change of coordinates formula [DS11, Proposition 2.1]. Let

$$Q = \frac{2}{\gamma} + \frac{\gamma}{2} \quad \text{for} \quad \gamma \in [0,2). \tag{2.9}$$

Suppose that D_1, D_2 are planar domains and $\varphi\colon D_1 \to D_2$ is a conformal map. If h_2 is (a form of) a GFF on D_2 and

$$h_1 = h_2 \circ \varphi + Q \log |\varphi'| \tag{2.10}$$

then

$$\mu_{h_1}(A) = \mu_{h_2}(\varphi(A)) \quad \text{and} \quad \nu_{h_1}(A) = \nu_{h_2}(\varphi(A)) \tag{2.11}$$

for all measurable sets A. This allows us to define an equivalence relation on pairs (D,h) by declaring (D_1,h_1) and (D_2,h_2) to be equivalent if h_1 and h_2 are related as in (2.10). An equivalence class of such a (D,h) is then referred to as a quantum surface.

More generally, suppose that D_1, D_2 are planar domains, $x_1^i, \ldots, x_n^i \in \overline{D}_i$ for $i = 1,2$ are given points, and h_i is a distribution on D_i. Then we say that the marked quantum surfaces $(D_i, h_i, x_1^i, \ldots, x_n^i)$ are equivalent if there exists a conformal transformation $\varphi\colon D_1 \to D_2$, $\varphi(x_j^1) = x_j^2$ for each $1 \le j \le n$, and h_1, h_2 are related as in (2.10).

In this work, we will be primarily interested in two types of quantum surfaces, namely quantum disks and spheres. We will remind the reader of the particular construction of a quantum sphere that we will be interested in for this work in Section 4.1. We also refer the reader to [MS15b] and [DMS14] for a careful definition of a quantum disk as well as several equivalent constructions of a quantum sphere.

2.3 Conformal removability

An LQG surface can be obtained by endowing a topological surface with both a good measure and a conformal structure in a random way. (And we can imagine that these two structures are added in either order.) Given two topological disks with boundary (each endowed with a good area measure in the interior, and a good length measure on the boundary) it is a simple matter to produce a new good-measure-endowed topological surface by taking a quotient that involves gluing (all or part of) the boundaries to each other in a boundary length preserving way.

The problem of *conformally welding* two surfaces is the problem of obtaining a conformal structure on the combined surface, given the conformal structure on the individual surfaces. (See, e.g., [Bis07] for further discussion and references.) To make sense of this idea, we will draw from the theory of *removable sets*, as explained below.

A compact subset K of a domain $D \subseteq \mathbf{C}$ is called (conformally) **removable** if every homeomorphism from D into $\mathbf{C}$ that is conformal on $D \setminus K$ is also conformal on all of D. A Jordan domain $D \subseteq \mathbf{C}$ is said to be a **Hölder domain** if any conformal transformation from $\mathbf{D}$ to D is Hölder continuous all of the way up to $\partial\mathbf{D}$. It was shown by Jones and Smirnov [JS00] that if $K \subseteq D$ is the boundary of a Hölder domain, then K is removable; it is also noted there that if a compact set K is removable as a subset of D, then it is removable in any domain containing K, including all of $\mathbf{C}$. Thus, at least for compact sets K, one can speak of removability without specifying a particular domain D.

The following proposition illustrates the importance of removability in the setting of quantum surfaces (see also [DMS14, Section 1.5]):

Proposition 2.1. *Suppose that (D, h) is a quantum surface, $K \subseteq \overline{D}$ is compact such that $D \setminus K = D_1 \cup D_2$ for D_1, D_2 disjoint. Suppose that (D', h') is another quantum surface, $K' \subseteq \overline{D}'$ is compact such that $D' \setminus K' = D_1' \cup D_1'$ for D_1', D_2' disjoint. Assume that (D_j, h) is equivalent to (D_j', h') as a quantum surface for $j = 1, 2$ and that, furthermore, there exists conformal transformations $\varphi_j \colon D_j \to D_j'$ for $j = 1, 2$ which extend to a homeomorphism $D \to D'$. If K is conformally removable, then (D, h) and (D', h') are equivalent as quantum surfaces.*

Proof. This follows immediately from the definition of conformal removability. □

One example of a setting in which Proposition 2.1 applies is when the quantum surface (D, h) is given by a so-called quantum wedge and K is the range of an SLE_κ curve η for $\kappa \in (0, 4)$ [She15, DMS14]. A quantum wedge naturally comes with two marked points x and y, which are also the seed and the target point of the SLE curve. In this case, the conformal maps φ_j (which are defined on the left and right components of $D \setminus K$) are chosen so that the quantum length of the image of a segment of η as

measured from the left and right sides matches up. With this choice, the φ_j extend to a homeomorphism of the whole domain and it shown in [RS05] that the range of η is almost surely conformally removable, so Proposition 2.1 applies.

When we apply Proposition 2.1 in the current work, we will not always be welding quantum surfaces according to quantum boundary length as in [She15, DMS14] because it will not always be obviously true that we will be in a setting in which ∂D_1 and ∂D_2 have an intrinsically defined quantum boundary length. This may *a priori* be the case for the QLE$(8/3, 0)$ process we construct here because we will not rule out the possibility that the process contains "spikes" (i.e., we will not show in this work that it is the case that the range of a QLE$(8/3, 0)$ is equal to the closure of its interior; this will be a consequence of the results of [MS16a]).

3 Eden model and percolation interface

In this section we briefly recall a few constructions from [MS13b, Section 2], together with some figures included there. Figure 3.1 shows a triangulation T of the sphere with two distinguished edges e_1 and e_2, and the caption describes a mechanism for choosing a random path in the dual graph of the triangulation, consisting of distinct triangles $t_1, t_2, \ldots, t_k$, that goes from e_1 to e_2. It will be useful to imagine that we begin with a single 2-gon and then grow the path dynamically, exploring new territory as we go. At any given step, we keep track of the total number edges on the boundary of the already-explored region and the number of vertices remaining to be seen in the component of the unexplored region that contains the target edge. The caption of Figure 3.2 explains one step of the exploration process. The exploration process induces a Markov chain on the set of pairs (m, n) with $m \geq 0$ and $n \geq 0$. In this chain, the n coordinate is almost surely non-increasing, and the m coordinate can only increase by 1 when the n coordinate decreases by 1.

Now consider the version of the Eden model in which new triangles are only added to the unexplored region containing the target edge, as illustrated Figure 3.3. In both Figure 3.1 and Figure 3.3, each time an exploration step separates the unexplored region into two pieces (each containing at least one triangle) we refer to the one that does not contain the target as a *bubble*. The exploration process described in Figure 3.1 created two bubbles (the two small white components), and the exploration process described in Figure 3.3 created one (colored blue). We can interpret the bubble as a triangulation of a polygon, rooted at a boundary edge (the edge it shares with the triangle that was observed when the bubble was created).

The specific growth pattern in Figure 3.3 is very different from the one depicted in Figure 3.1. However, the analysis used in Figure 3.2 applies equally well to both scenarios. The only difference between the two is that in Figure 3.3 one re-randomizes the seed edge (choosing it uniformly from all possible values) after each step.



Figure 3.1: **Upper left:** a triangulation of the sphere together with two distinguished edges colored green. **Upper right:** It is conceptually useful to "fatten" each green edge into a 2-gon. We fix a distinguished non-self-intersecting dual-lattice path p (dotted red line) from one 2-gon to the other. **Bottom:** Vertices are colored red or blue with i.i.d. fair coins. There is then a unique dual-lattice path from one 2-gon to the other (triangles in the path colored orange) such that each edge it crosses either has opposite-colored endpoints and *does not* cross p, or has same-colored endpoints and *does* cross p. The law of the orange path does not depend on the choice of p, since shifting p across a vertex has the same effect as flipping the color of that vertex.

In either of these models, we can define C_k to be the boundary of the target-containing unexplored region after k steps. If (M_k, N_k) is the corresponding Markov chain, then the length of C_k is $M_k + 2$ for each k. Let D_k denote the union of the edges and vertices in C_k, the edges and vertices in C_{k-1} and the triangle and bubble (if applicable) added at step k, as in Figure 3.4. We refer to each D_k as a *necklace* since it typically contains a cycle of edges together with a cluster of one or more triangles hanging off of it. The analysis used in Figure 3.2 (and discussed above) immediately implies the following:

Proposition 3.1. *Consider a random rooted triangulation of the sphere with a fixed*



Figure 3.2: When "exploring" the polygon adjacent to a chosen edge on the boundary of the unexplored region, one encounters either a triangle whose third vertex is in the interior (leftmost two figures), a triangle whose third vertex is on the boundary (next two figures) or the terminal green 2-gon (right figure).



Figure 3.3: Same as Figure 3.1 except that one explores using the Eden model instead of percolation. At each step, one chooses a uniformly random edge on the boundary of the unexplored region containing the target and explores the face incident to that edge. The faces are numbered according to the order in which they were explored. When the unexplored region is divided into two pieces, each with one or more triangles, the piece without the target is called a *bubble* and is never subsequently explored by this process. In this figure there is only one bubble, which is colored blue.

number $n > 2$ of vertices together with two distinguished edges chosen uniformly from the set of possible edges. If we start at one edge and explore using the Eden model as in Figure 3.3, or if we explore using the percolation interface of Figure 3.1, we will find that the following are the same:

1. *The law of the Markov chain (M_k, N_k) (which terminates when the target 2-gon is reached).*

2. *The law of the total number of triangles observed before the target is reached.*

3. *The law of the sequence D_k of necklaces.*

Indeed, one way to construct an instance of the Eden model process is to start with an instance of the percolation interface exploration process and then randomly rotate the necklaces in the manner illustrated in Figure 3.4.



Figure 3.4: **Left:** the first four necklaces (separated by white space) generated by an Eden model exploration. **Middle:** one possible way of identifying the vertices on the outside of each necklace with those on the inside of the next necklace outward. **Right:** The map with exploration associated to this identification. If a necklace has n vertices on its outer boundary, then there are n ways to glue this outer boundary to the inner boundary of the next necklace outward. It is natural to choose one of these ways uniformly at random, independently for each consecutive pair of necklaces. Intuitively, we imagine that before gluing them together, we randomly spin the necklaces like the reels of a slot machine. A fanciful interpretation of Proposition 3.1 is that if we take a percolation interface exploration as in Figure 3.1 (which describes a sequence of necklaces) and we pull a slot machine lever that spins the necklaces independently (see [MS13b]), then we end up with an Eden model exploration of the type shown in Figure 3.3.

4 Infinite measures on quantum spheres

4.1 Lévy excursion description of doubly-marked quantum spheres

The purpose of this section is to review the results established in [MS15b] which are relevant for this article. First, we let $\mathsf{M}^2_{\mathrm{SPH}}$ be the measure which is defined as follows. Suppose that X_t is a 3/2-stable Lévy process with only upward jumps and let I_t be its running infimum. Then we let N be the Itô excursion measure associated with the excursions that $X_t - I_t$ makes from 0. The law of the duration of such an excursion follows a power law. Indeed, following [Ber96], the process of sampling N can be described as follows (see [Ber96, Section VIII.4]):

1. Pick a lifetime t from the measure $ct^{\rho-2}dt = ct^{-5/3}dt$ on $\mathbf{R}_+$ where dt denotes Lebesgue measure and $c > 0$ is a constant. Here, $\rho = 1 - 1/\alpha = 1/3$ (where $\alpha = 3/2$) is the so-called positivity parameter of the process [Ber96, Section VIII.1].
2. Given t, pick an excursion of length from the normalized excursion measure $\mathbf{n}$ associated with a 3/2-stable Lévy process with only positive jumps and then rescale it spatially and in time so that it has length t.

As explained in [MS15b], we can construct a doubly-marked quantum sphere $(\mathcal{S}, x, y)$ decorated by a non-crossing path η' connecting x and y from such an excursion $e\colon [0,T] \to \mathbf{R}_+$ as follows. For each upward jump of e, we sample a conditionally independent quantum disk whose boundary length is equal to the size of the jump. We assume that each of the quantum disks has a marked boundary point sampled uniformly from its boundary measure together with a uniformly chosen orientation of its boundary. We assume that the marked points and orientations are chosen conditionally independently given the realizations of the quantum disks. Let $\mathcal{D}$ denote the collection of marked and oriented quantum disks sampled in this way. Then the pair $e, \mathcal{D}$ together uniquely determines a doubly-marked surface $(\mathcal{S}, x, y)$ which is homeomorphic to the sphere together with a non-crossing path η' which connects x and y. In particular, the time-reversal of e describes the evolution of the quantum boundary length of the complementary component of $\eta'([0,t])$ in $\mathcal{S}$ which contains y and the jumps of e describe the boundary lengths of the quantum disks that η' cuts off from y. For each $t \geq 0$, we let K_t be the component of $\mathcal{S} \setminus \eta'([0,t])$ which contains y. The time-parameterization of η' so that the quantum boundary length of the component of $\mathcal{S} \setminus K_t$ which contains y is equal to $e(T-t)$ is the so-called **quantum natural time** introduced in [DMS14].

One of the main results of [MS15b] is that a doubly-marked surface/path pair $(\mathcal{S}, x, y), \eta'$ produced from $\mathsf{M}^2_{\mathrm{SPH}}$ *conditioned* to have quantum mass equal to 1 (although $\mathsf{M}^2_{\mathrm{SPH}}$ is infinite, this conditioning yields a probability measure) has the law of the unit area quantum sphere constructed in [DMS14], the points x, y conditional on $\mathcal{S}$ are chosen

uniformly at random from the quantum measure, and the conditional law of η' given $x, y, \mathcal{S}$ is that of a whole-plane SLE_6 process connecting x and y. This holds more generally when we condition the surface to have quantum mass equal to m for any fixed $m > 0$ except in this setting $\mathcal{S}$ is a quantum sphere of mass m rather than 1. (As noted earlier, a sample from the law of such a surface can be produced by starting with a unit area quantum sphere and then scaling its associated mass measure by the factor m.)

This relationship between $\mathsf{M}^2_{\mathrm{SPH}}$ and the law of a unit area quantum sphere decorated with an independent whole-plane SLE_6 process implies that $\mathsf{M}^2_{\mathrm{SPH}}$ possesses certain symmetries. These symmetries will be important later on so we will pause for a moment to point them out.

- If we condition on $\mathcal{S}$, then the points x and y are both chosen independently from the quantum area measure on $\mathcal{S}$.
- If we condition on x, y, and $\mathcal{S}$, then η' is whole-plane SLE_6 from x to y.
- The amount of quantum natural time elapsed for η' to travel from x to y is equal to T (the time corresponding to the Lévy excursion).

This also implies that $\mathsf{M}^2_{\mathrm{SPH}}$ is invariant under the operation of swapping x and y and then reversing the time of η' [MS13a] (with the quantum natural time parameterization). To see the symmetry of the quantum natural time parameterization under time-reversal, we have from the previous observations that the law of the ordered collection of bubbles cut off by η' from its target point is invariant under the operation of swapping x and y and reversing the time of η'. The claim thus follows because the quantum natural time parameterization can be constructed by fixing $j \in \mathbf{N}$, counting the number $N(e^{-j-1}, e^{-j})$ of bubbles cut off by η' with quantum boundary length in $[e^{-j-1}, e^{-j}]$, and then normalizing by a constant times the factor $e^{3/2j}$. That this is the correct normalization follows since the Lévy measure for a 3/2-stable Lévy process is given by a constant times $u^{-5/2}du$ where du denotes Lebesgue measure on $\mathbf{R}_+$. See, for example, [MS15b, Section 6.2] for additional discussion of this point as well as Remark 6.4 below in the context of the construction of QLE$(8/3, 0)$.

We also emphasize that under $\mathsf{M}^2_{\mathrm{SPH}}$, we have that:

- $\eta'(t)$ is distributed uniformly from the quantum boundary measure associated with the quantum surface parameterized by $\mathcal{S} \setminus K_t$ (see [MS15b, Proposition 6.4]).
- The components of $\mathcal{S} \setminus \eta'([0,t])$, viewed as quantum surfaces, are conditionally independent given their boundary lengths. Those components which do not contain y are quantum disks given their boundary lengths. The component which does contain y has the law of a quantum disk weighted by its total quantum area.

4.2 Weighted measures

Throughout, we will work with the following two measures which are defined with $\mathsf{M}^2_{\mathrm{SPH}}$ as the starting point. Namely, with T equal to the length of the associated Lévy excursion and X_t the quantum boundary length of the complementary component of $\eta'([0,t])$ containing y we write

$$d\mathsf{M}^2_{\mathrm{SPH,W}} = \mathbf{1}_{[0,T]}(\mathfrak{t})d\mathfrak{t}d\mathsf{M}^2_{\mathrm{SPH}} \quad \text{and} \tag{4.1}$$

$$d\mathsf{M}^2_{\mathrm{SPH,D}} = \mathbf{1}_{[0,T]}(\mathfrak{t})\frac{1}{X_{\mathfrak{t}}}d\mathfrak{t}d\mathsf{M}^2_{\mathrm{SPH}} = \frac{1}{X_{\mathfrak{t}}}d\mathsf{M}^2_{\mathrm{SPH,W}} \tag{4.2}$$

where $d\mathfrak{t}$ denotes Lebesgue measure. We note that the marginal of $\mathsf{M}^2_{\mathrm{SPH,W}}$ on $(\mathcal{S}, x, y)$ and η' is given by $Td\mathsf{M}^2_{\mathrm{SPH}}$, i.e. by weighting $\mathsf{M}^2_{\mathrm{SPH}}$ by the length of the Lévy excursion. (The additional subscript "W" is to indicate that $\mathsf{M}^2_{\mathrm{SPH,W}}$ is a weighted measure.) It will be convenient throughout to think of $\mathsf{M}^2_{\mathrm{SPH,W}}$ as a measure on triples $\left(\mathfrak{t},\, (\mathcal{S}, x, y),\, \eta'\right)$ where $\mathfrak{t}$ is a uniformly random point chosen from the total length of the Lévy excursion.

The marginal of $\mathsf{M}^2_{\mathrm{SPH,D}}$ on $(\mathcal{S}, x, y)$ and η' is given by $Dd\mathsf{M}^2_{\mathrm{SPH}}$ where

$$D = \int_0^T X_s^{-1} ds.$$

As we will see later, after the QLE "reshuffling" this will have the interpretation of taking $\mathsf{M}^2_{\mathrm{SPH}}$ and then weighting it by the amount of QLE$(8/3, 0)$ quantum distance from x to y. (The additional subscript "D" is to indicate that $\mathsf{M}^2_{\mathrm{SPH,D}}$ is the "distance weighted" measure.)

We finish this section by recording the following proposition which relates the conditional law of $\mathsf{M}^2_{\mathrm{SPH,W}}$ and $\mathsf{M}^2_{\mathrm{SPH,D}}$ given $\mathfrak{t}$ and $X_{\mathfrak{t}}$ to $\mathsf{M}^2_{\mathrm{SPH}}$.

Proposition 4.1. *(i) Given $\mathfrak{t}$, the conditional distribution of $\mathsf{M}^2_{\mathrm{SPH,W}}$ is the same as the conditional distribution of $\mathsf{M}^2_{\mathrm{SPH}}$ when we condition on the event that the length of the Lévy excursion is at least $\mathfrak{t}$.*

(ii) For both $\mathsf{m} = \mathsf{M}^2_{\mathrm{SPH,W}}$ and $\mathsf{m} = \mathsf{M}^2_{\mathrm{SPH,D}}$, given $\mathfrak{t}$ and $X_{\mathfrak{t}}$, the conditional distribution of m is the same as the conditional distribution of $\mathsf{M}^2_{\mathrm{SPH}}$ when we condition on the event that the length of the Lévy excursion is at least $\mathfrak{t}$ and the given value of $X_{\mathfrak{t}}$.

Proof. We will explain the argument in the case that $\mathsf{m} = \mathsf{M}^2_{\mathrm{SPH,W}}$; the same argument gives part (ii).

If we fix the value of $\mathfrak{t}$, the conditional distribution of the Lévy excursion in the definition of $\mathsf{M}^2_{\mathrm{SPH,W}}$ is given by $c\mathbf{1}_{[\mathfrak{t},\infty)}(T)d\mathbf{n}_T T^{-5/3}dT$ where $\mathbf{n}_T$ is the measure on 3/2-stable Lévy excursions which arises by scaling $\mathbf{n}$ spatially and in time so that the excursion length is equal to T. This representation clearly implies (i). A similar argument gives (ii). □

4.3 Continuum scaling exponents

We now determine the distribution of D and A under $\mathsf{M}^2_{\mathrm{SPH}}$.

Proposition 4.2. *There exists constants $c_0, c_1 > 0$ such that*

$$\mathsf{M}^2_{\mathrm{SPH}}[D \geq t] = \frac{c_0}{t^2} \quad \textit{and} \tag{4.3}$$

$$\mathsf{M}^2_{\mathrm{SPH}}[A \geq a] = \frac{c_1}{a^{1/2}}. \tag{4.4}$$

Note that the exponents in (4.3), (4.4) match the corresponding exponents derived in [MS15a].

Proof of Proposition 4.2. We note that (4.4) is explained just after the statement of [MS15b, Theorem 1.4]. Therefore we only need to prove (4.3).

For an excursion $e\colon [0,T] \to \mathbf{R}_+$ sampled from N we write $e^* = \sup_{t\in[0,T]} e(t)$. By scaling and the explicit form of N described above, it is not difficult to see by making the change of variables $u = tT^{-2/3}$ that there exists a constant $c_0 > 0$ such that

$$\mathsf{N}[e^* \geq t] = \int_0^\infty \mathbf{n}[e^* \geq tT^{-2/3}]T^{-5/3}dT = \frac{c_0}{t}. \tag{4.5}$$

For each $\epsilon > 0$, we let $T_\epsilon = \inf\{t \geq 0 : e(t) \geq \epsilon\}$ and let

$$D^\epsilon = \int_{T_\epsilon}^T \frac{1}{X_s} ds.$$

Suppose that Y is a $u^{3/2}$-CSBP with $Y_0 = \epsilon$. Let $\zeta = \inf\{t \geq 0 : Y_t = 0\}$ be the extinction time of Y. Then it follows from (2.1) and (2.8) that

$$\mathbf{P}[\zeta \leq t] = \lim_{\lambda\to\infty} \mathbf{E}[e^{-\lambda Y_t}] = \lim_{\lambda\to\infty} e^{\epsilon u_t(\lambda)} = e^{-4\epsilon/t^2}.$$

Therefore

$$\mathbf{P}[\zeta \geq t] = \frac{4\epsilon}{t^2} + o(\epsilon) \quad \text{as} \quad \epsilon \to 0.$$

By combining this with (2.6), it therefore follows that there exists a constant $c_1 > 0$ such that

$$\begin{aligned}\mathsf{M}^2_{\mathrm{SPH}}[D^\epsilon \geq t] &= \mathsf{M}^2_{\mathrm{SPH}}[D^\epsilon \geq t \,|\, e^* \geq \epsilon]\mathsf{M}^2_{\mathrm{SPH}}[e^* \geq \epsilon] \\ &= \frac{c_1\epsilon}{t^2} \cdot \frac{c_0}{\epsilon} + o(1) = \frac{c_0c_1}{t^2} + o(1) \quad \text{as} \quad \epsilon \to 0.\end{aligned}$$

Sending $\epsilon \to 0$ implies the result. □

5 Meeting in the middle

In this section, we shall assume that we are working in the setting described in Section 4.2. The main result is the following theorem which proves a certain symmetry statement for the measure $\mathsf{M}^2_{\mathrm{SPH,D}}$. This result will later be used in Section 7 to prove an analogous symmetry result for QLE$(8/3, 0)$ which, in turn, is one of the main inputs in the proof of Theorem 1.1. See Figure 5.1 for an illustration of the result.



Figure 5.1: Illustration of the setup for Theorem 5.1, the main result of Section 5. Shown is a doubly-marked quantum sphere $(\mathcal{S}, x, y)$ decorated with a whole-plane SLE_6 process η' connecting x and y. The blue path shows η' drawn up to a time $\mathfrak{t}$ which is uniform in the total amount of quantum natural time required by η' to connect x and y and the red path is the time-reversal $\overline{\eta}'$ drawn up until the first time $\bar{\mathfrak{t}}$ that it hits $\eta'([0,\mathfrak{t}])$. Let $\mathcal{X}$ (resp. $\overline{\mathcal{X}}$) consist of $\eta'|_{[0,\mathfrak{t}]}$ (resp. $\overline{\eta}'|_{[0,\bar{\mathfrak{t}}]}$) and the part of $\mathcal{S}$ separated by $\eta'([0,\mathfrak{t}])$ (resp. $\overline{\eta}'([0,\bar{\mathfrak{t}}])$) from y (resp. x) and let $\mathcal{B}$ be the part of $\mathcal{S}$ which is not in $\mathcal{X}$ and $\overline{\mathcal{X}}$. In the illustration, $\mathcal{X}$ (resp. $\overline{\mathcal{X}}$) is shown in light blue (resp. light red). In Theorem 5.1, we show that the $\mathsf{M}^2_{\mathrm{SPH,D}}$ distribution of the triple $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$ is invariant under the operation of swapping $\mathcal{X}$ and $\overline{\mathcal{X}}$.

Theorem 5.1. *Suppose that $(\mathcal{S}, x, y)$ is a doubly-marked quantum sphere, η' is a path on $\mathcal{S}$ from x to y, and $\mathfrak{t} \geq 0$, where these objects are sampled from $\mathsf{M}^2_{\mathrm{SPH,D}}$ as in Section 4.2. Let $\overline{\eta}'$ be the time-reversal of η' and let $\bar{\mathfrak{t}}$ be the first time that $\overline{\eta}'$ hits $\eta'([0,\mathfrak{t}])$. Let*

- *$\mathcal{X}$ be the path-decorated and beaded quantum surface parameterized by the union of $\eta'|_{[0,\mathfrak{t}]}$ and the part of $\mathcal{S}$ separated from y by $\eta'([0,\mathfrak{t}])$,*
- *$\overline{\mathcal{X}}$ be the path-decorated and beaded quantum surface parameterized by the union of $\overline{\eta}'|_{[0,\bar{\mathfrak{t}}]}$ and the part of $\mathcal{S}$ separated from x by $\overline{\eta}'([0,\bar{\mathfrak{t}}])$, and*
- *$\mathcal{B}$ be the (disk-homeomorphic) quantum surface parameterized by $\mathcal{S} \setminus (\mathcal{X} \cap \overline{\mathcal{X}})$.*

We view $\mathcal{X}$ (resp. $\overline{\mathcal{X}}$) as a random variable taking values in the space of path-decorated and beaded quantum surfaces with a marked point which corresponds to $\eta'(\mathfrak{t})$ (resp. $\overline{\eta}'(\overline{\mathfrak{t}})$) and we view $\mathcal{B}$ as a random variable taking values in the space of quantum surfaces. Under $\mathsf{M}^2_{\mathrm{SPH,D}}$, we have that $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ are conditionally independent given the quantum lengths of the boundaries of $\mathcal{X}$ and $\overline{\mathcal{X}}$ (which together determine the boundary length of $\mathcal{B}$). Moreover, the $\mathsf{M}^2_{\mathrm{SPH,D}}$ distribution of $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$ is invariant under the operation of swapping $\mathcal{X}$ and $\overline{\mathcal{X}}$.

Remark 5.2. We emphasize that the assertion of Theorem 5.1 does not give that the surface $\mathcal{S}$ decorated by the paths $\eta'|_{[0,\mathfrak{t}]}$ and $\overline{\eta}'|_{[0,\overline{\mathfrak{t}}]}$ is invariant under swapping $\eta'|_{[0,\mathfrak{t}]}$ and $\overline{\eta}'|_{[0,\overline{\mathfrak{t}}]}$. This statement does not hold because there is an asymmetry in that $\eta'(\mathfrak{t}) \notin \overline{\eta}'([0,\overline{\mathfrak{t}}])$ while we have that $\overline{\eta}'(\overline{\mathfrak{t}}) \in \eta'([0,\mathfrak{t}])$. Rather, Theorem 5.1 implies that the induced distribution on triples of $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$ is invariant under swapping $\mathcal{X}$ and $\overline{\mathcal{X}}$. The difference between the two statements is that the first statement depends on how the surfaces which correspond to $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ are glued together (which determines the locations of the tips) while the second statement does not depend on how everything is glued together. In the QLE$(8/3, 0)$ analog of Theorem 5.1, which is stated as Theorem 7.1 below (and will be derived as a consequence of Theorem 5.1), we do have the symmetry of the whole picture because the tips are "lost" in the "reshuffling" procedure used to construct QLE$(8/3, 0)$ from SLE_6.

Our ultimate aim in this section is to deduce Theorem 5.1 by showing that the $\mathsf{M}^2_{\mathrm{SPH,D}}$ distribution on $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$ can be constructed from $\mathsf{M}^2_{\mathrm{SPH,W}}$ by "conditioning" on the event that $\mathfrak{t}$ is a cut time for η'. This will indeed lead to the desired result because, as we will explain in Proposition 5.3 just below, $\mathsf{M}^2_{\mathrm{SPH,W}}$ possesses a symmetry property which is similar to that described in Theorem 5.1.

Proposition 5.3. *Suppose that $(\mathcal{S}, x, y)$ is a doubly-marked quantum sphere, η' is a path connecting x to y, and $\mathfrak{t} \geq 0$, where these objects are sampled from $\mathsf{M}^2_{\mathrm{SPH,W}}$ as defined in Section 4.2. Let $\overline{\tau}$ be the first time that $\overline{\eta}'$ hits $\eta'(\mathfrak{t})$. Let*

- *$\mathcal{Y}$ be the path-decorated and beaded quantum surface parameterized by the union of $\eta'|_{[0,\mathfrak{t}]}$ and the part of $\mathcal{S}$ which is separated from y by $\eta'([0,\mathfrak{t}])$,*
- *$\overline{\mathcal{Y}}$ be the path-decorated and beaded quantum surface parameterized by the union of $\overline{\eta}'|_{[0,\overline{\tau}]}$ and the part of $\mathcal{S}$ which is separated from x by $\overline{\eta}'([0,\overline{\tau}])$, and*
- *$\mathcal{Q}$ be the quantum surface parameterized by $\mathcal{S} \setminus (\mathcal{Y} \cup \overline{\mathcal{Y}})$.*

We view $\mathcal{Y}$ (resp. $\overline{\mathcal{Y}}$) as a random variable taking values in the space of path-decorated and beaded quantum surfaces with a single marked point which corresponds to $\eta'(\mathfrak{t})$ (resp. $\overline{\eta}'(\overline{\tau})$) and we view $\mathcal{Q}$ as a random variable taking values in the space of quantum surfaces. Then $\mathsf{M}^2_{\mathrm{SPH,W}}$ is invariant under the operation of swapping $\mathcal{Y}$ and $\overline{\mathcal{Y}}$. Moreover, under $\mathsf{M}^2_{\mathrm{SPH,W}}$, we have that $\mathcal{Y}$, $\overline{\mathcal{Y}}$, and $\mathcal{Q}$ are conditionally independent given the quantum



Figure 5.2: Illustration of the setup for Proposition 5.3 (continuation of Figure 5.1). Let $\mathcal{Y}$ (resp. $\overline{\mathcal{Y}}$) consist of $\eta'|_{[0,\mathfrak{t}]}$ (resp. $\overline{\eta}'|_{[0,\overline{\tau}]}$ where $\overline{\tau}$ is the first time that $\overline{\eta}'$ hits $\eta'(\mathfrak{t})$) and the part of $\mathcal{S}$ separated by $\eta'([0,\mathfrak{t}])$ (resp. $\overline{\eta}'([0,\overline{\tau}])$) from y (resp. x) and let $\mathcal{Q}$ be the part of $\mathcal{S}$ which is not in $\mathcal{Y}$ and $\overline{\mathcal{Y}}$. In the illustration, $\mathcal{Y}$ (resp. $\overline{\mathcal{Y}}$) is shown in light blue (resp. light red and light green). In Proposition 5.3, we show that the $\mathsf{M}^2_{\mathrm{SPH,W}}$ distribution of the triple $(\mathcal{Y}, \overline{\mathcal{Y}}, \mathcal{Q})$ is invariant under the operation of swapping $\mathcal{Y}$ and $\overline{\mathcal{Y}}$.

lengths of the boundaries of $\mathcal{Y}$ and $\overline{\mathcal{Y}}$ (which together determine the boundary length of $\mathcal{Q}$).

Proof. We start with the first assertion of the proposition. As mentioned in Section 4.1, we know that if we condition on the quantum area m of $\mathcal{S}$ then the joint law of $(\mathcal{S}, x, y)$ and η' is given by a quantum sphere of quantum area m, x and y are independently and uniformly chosen from the quantum area measure, and, given x, y, η' is an independent whole-plane SLE_6 connecting x and y. By the reversibility of whole-plane SLE_6 established in [MS13a], it thus follows that the joint law of $(\mathcal{S}, x, y)$ and η' is invariant under swapping x and y and reversing the time of η' when m is fixed. Since $\mathfrak{t}$ is uniform in the amount of quantum natural time T required by η' to connect x and y, we have by symmetry that $\overline{\tau}$ is uniform in $[0, T]$. This proves the first part.

It follows from the construction of $\mathsf{M}^2_{\mathrm{SPH,W}}$ that $\mathcal{Y}$ is independent of $\overline{\mathcal{Y}}$ and $\mathcal{Q}$ given its quantum boundary length (recall Proposition 4.1) and, by symmetry, that $\overline{\mathcal{Y}}$ is independent of $\mathcal{Y}$ and $\mathcal{Q}$ given its boundary length. This implies the second assertion of the proposition. □

We will establish Theorem 5.1 using the following strategy. Suppose that we have a triple T, $(\mathcal{S}, x, y)$, η' which consists of a number $T \geq 0$, a doubly-marked finite volume quantum surface $\mathcal{S}$ homeomorphic to $\mathbf{S}^2$, and a non-crossing path η' on $\mathcal{S}$ connecting x and y such that T is equal to the total amount of quantum natural time taken by

η' to go from x to y. We assume that $\mathcal{S}$ is parameterized by $\mathbf{C}$ with 0 corresponding to x and ∞ corresponding to y. Let $\mathfrak{t}$ be uniform in $[0,T]$ and let U be the unbounded component of $\mathbf{C} \setminus \eta'([0,\mathfrak{t}])$. Let $\overline{\eta}'$ be the time-reversal of η' and let $\overline{\mathfrak{t}}$ be the first time that $\overline{\eta}'$ hits $\eta'([0,\mathfrak{t}])$. Fix $\epsilon > 0$ and let E_ϵ be the event that $\overline{\eta}'(\overline{\mathfrak{t}})$ is contained in the interval of ∂U starting from $\eta'(\mathfrak{t})$ and continuing in the counterclockwise direction until reaching quantum length ϵ. (In the case that ∂U has quantum length at most ϵ, we take this interval to be all of ∂U.) In other words, E_ϵ is the event that $\overline{\eta}'(\overline{\mathfrak{t}})$ is contained in the quantum length ϵ interval on the outer boundary of $\eta'([0,\mathfrak{t}])$ which is immediately to the left of $\eta'(\mathfrak{t})$. Let $\overline{\tau}$ be the first time that $\overline{\eta}'$ hits $\eta'(\mathfrak{t})$ and let F_ϵ be the event that $\overline{\eta}'([0,\overline{\tau}]) \cap \eta'([0,\mathfrak{t}])$ is contained in both the interval of ∂U centered at $\eta'(\mathfrak{t})$ with quantum length 2ϵ and the similarly defined interval on the outer boundary of $\overline{\eta}'([0,\overline{\tau}])$ with the roles of η' and $\overline{\eta}'$ swapped.

The main two steps in the proof of Theorem 5.1 are to show that (with $\mathsf{M}^2_{\mathrm{SPH,W}}$ viewed as a measure on $(\mathcal{Y}, \overline{\mathcal{Y}}, \mathcal{Q})$ and $\mathsf{M}^2_{\mathrm{SPH,D}}$ viewed as a measure on $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$) we have

$$\epsilon^{-1}\mathbf{1}_{E_\epsilon} d\mathsf{M}^2_{\mathrm{SPH,W}} \to d\mathsf{M}^2_{\mathrm{SPH,D}} \quad \text{as} \quad \epsilon \to 0 \quad \text{and} \tag{5.1}$$

$$\epsilon^{-1}\mathbf{1}_{F_\epsilon} d\mathsf{M}^2_{\mathrm{SPH,W}} \to c_0 d\mathsf{M}^2_{\mathrm{SPH,D}} \quad \text{as} \quad \epsilon \to 0, \tag{5.2}$$

where $c_0 > 0$ is a fixed constant. We will prove (5.1) as a step in proving (5.2) and deduce (5.2) from (5.1) by showing that the conditional probability of F_ϵ given E_ϵ converges as $\epsilon \to 0$ to the positive constant c_0. The notion of convergence is given by weak convergence with respect to a topology that we will introduce in the proof of Theorem 5.1. (The exact choice of topology is not important as long as it generates the full σ-algebra.) The result then follows because, for each $\epsilon > 0$, $\epsilon^{-1}\mathbf{1}_{F_\epsilon} d\mathsf{M}^2_{\mathrm{SPH,W}}$ is invariant under the operation of swapping $\mathcal{Y}$ and $\overline{\mathcal{Y}}$ since $\mathsf{M}^2_{\mathrm{SPH,W}}$ itself is invariant under this operation and F_ϵ is defined in a manner which is symmetric in $\eta'|_{[0,\mathfrak{t}]}$ and $\overline{\eta}'|_{[0,\overline{\tau}]}$. (We note that $\overline{\tau} \to \overline{\mathfrak{t}}$ as $\epsilon \to 0$.)

Throughout the remainder of this section, we will write $\mathbf{P}_u$ for the probability which gives the conditional law of η' and the remaining surface given $X_{\mathfrak{t}} = u$ under either $\mathsf{M}^2_{\mathrm{SPH,W}}$ or $\mathsf{M}^2_{\mathrm{SPH,D}}$. (As we pointed out in Proposition 4.1, this conditional law is the same under both $\mathsf{M}^2_{\mathrm{SPH,W}}$ and $\mathsf{M}^2_{\mathrm{SPH,D}}$.)

Lemma 5.4. *We have that*

$$\mathbf{P}_u[E_\epsilon] = \frac{\epsilon}{u} \wedge 1.$$

Proof. This follows because the location of $\eta'(\mathfrak{t})$ on $\partial\mathcal{Y}$ is uniform from the quantum measure given the quantum boundary length of $\partial\mathcal{Y}$. □

In our next lemma, we show that the conditional law of $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$ given $(\mathfrak{t}, X_{\mathfrak{t}})$ sampled from $\mathsf{M}^2_{\mathrm{SPH,W}}$ does not change when we further condition on E_ϵ. We emphasize, however, that the conditional law of the part of η' which connects $\eta'(\mathfrak{t})$ and $\overline{\eta}'(\overline{\mathfrak{t}})$ (the green path segment in Figure 5.2) given E_ϵ is not the same as its unconditioned law.

Lemma 5.5. *Suppose that $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ are as in the statement of Theorem 5.1. The joint law of $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ under $\mathsf{M}^2_{\mathrm{SPH}}$ conditional on $(\mathfrak{t}, X_{\mathfrak{t}})$ is equal to the joint law of $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ under $\mathsf{M}^2_{\mathrm{SPH}}$ conditional on $(\mathfrak{t}, X_{\mathfrak{t}})$ and E_ϵ. The same holds with either $\mathsf{M}^2_{\mathrm{SPH,W}}$ or $\mathsf{M}^2_{\mathrm{SPH,D}}$ in place of $\mathsf{M}^2_{\mathrm{SPH}}$.*

Proof. This follows from the same argument used to prove Lemma 5.4. □

In our next lemma, we show that the local behavior of $X_{\mathfrak{t}}$ near $\mathfrak{t}$ is that of a 3/2-stable Lévy process with only downward jumps.

Lemma 5.6. *Suppose that Y is given by the time-reversal of a 3/2-stable Lévy excursion with only upward jumps of length at least t conditioned so that $Y_t = u$. Let $\mathcal{Z}$ be the Radon-Nikodym derivative between the law of $s \mapsto Y_{t+s}$ for $s \in [0, \epsilon]$ with respect to the law of a 3/2-stable Lévy process with only downward jumps in $[0, \epsilon]$ starting from u. Then we have that $\mathcal{Z} \to 1$ in probability as $\epsilon \to 0$.*

Proof. This follows, for example, from [MS15a, Lemma 3.19]. □

We will now describe the local behavior of a surface sampled from $\mathbf{P}_u$ near $\eta'(\mathfrak{t})$ both conditioned on E_ϵ and unconditioned. See Figure 5.3 for an illustration of the setup.

Proposition 5.7. *Fix $u, \epsilon > 0$ and let U be the unbounded component of $\mathbf{C} \setminus \eta'([0,t])$. Let $\varphi_1 \colon U \to \mathbf{H}$ be the unique conformal map which takes ∞ to i and $\eta'(t)$ to 0 and let $\varphi_2 \colon \mathbf{H} \to \mathbf{H}$ be the conformal map which corresponds to scaling so that with $\varphi = \varphi_2 \circ \varphi_1$ the quantum length assigned to $[-1, 0]$ by the quantum boundary measure associated with the field $\widetilde{h} = h \circ \varphi^{-1} + Q \log |(\varphi^{-1})'| + \frac{2}{\gamma} \log \epsilon^{-1}$ is equal to 1. Let $\widetilde{\eta}' = \varphi(\eta')$.*

Let $\overline{\sigma}$ be any stopping time for the filtration $\mathcal{F}_s$ generated by the time-reversal of η' and $\eta'|_{[0,t]}$, which almost surely occurs before the path first hits $\eta'([0,t])$. We have that the joint law of the quantum surface $(\mathbf{H}, \widetilde{h})$ and path $\widetilde{\eta}'$ under $\mathbf{P}_u[\cdot \,|\, E_\epsilon, \mathcal{F}_{\overline{\sigma}}]$ converges as $\epsilon \to 0$ to a pair consisting of a $\sqrt{8/3}$-quantum wedge with scaling factor chosen so that the quantum boundary length of $[-1, 0]$ is equal to 1 and an independent path $\widetilde{\eta}'$ whose law can be sampled from using the following steps.

- *Sample $w \in [-1, 0]$ according to Lebesgue measure*
- *Sample an $\mathrm{SLE}_6(2;2)$ process $\widetilde{\eta}'_w$ from w to ∞ with force points located at w^-, w^+; let V_0 be the component of $\mathbf{H} \setminus \widetilde{\eta}'_w$ with 0 on its boundary*
- *Given $\widetilde{\eta}'_w$, sample an SLE_6 process $\widetilde{\eta}'_0$ in V_0 from 0 to w*
- *Take the concatenation of $\widetilde{\eta}'_0$ and $\widetilde{\eta}'_w$.*



Figure 5.3: Illustration of the setup of Proposition 5.7. Shown on the left in blue is a path η' on top of a doubly-marked quantum surface parameterized by $(\mathbf{C}, h)$ drawn up to time t and in red the time-reversal $\overline{\eta}'$ of η' up to a time $\overline{\sigma}$ before it hits $\eta'([0,t])$. We show in Proposition 5.7 that under $\mathbf{P}_u$ conditioned on $\overline{\eta}'|_{[0,\overline{\sigma}]}$ and on the event that $\overline{\eta}'$ first hits $\eta'(t)$ in the counterclockwise segment of the outer boundary of $\eta'([0,t])$ of length ϵ the local behavior of the surface near $\eta'(t)$ is described by a $\sqrt{8/3}$-quantum wedge and the local behavior η' is described by an independent path which can be constructed by concatenating two SLE_6-type curves.

In the statement of Proposition 5.7, we emphasize that since $\frac{\kappa'}{2} - 2$ is equal to 1 for $\kappa' = 6$, we have that $\widetilde{\eta}'_w$ almost surely does not hit $\mathbf{R}$ except at its starting point; see [MS12a, Remark 2.3]. Before we give the proof of Proposition 5.7, we will need to collect several intermediate results.

Lemma 5.8. *Suppose that η' is a chordal SLE_6 process from 0 to ∞ in $\mathbf{H}$ and let $\overline{\eta}'$ be its time-reversal. Let $[x,y] \subseteq \mathbf{R}$ be an interval containing a neighborhood of 0, let $\overline{\tau}$ be the first time t that $\overline{\eta}'(t) \in [x,y]$, and let $w = \overline{\eta}'(\overline{\tau})$. Let τ be the last time that η' hits w (so that $\eta'|_{[\tau,\infty)}$ is the time-reversal of $\overline{\eta}'|_{(0,\overline{\tau}]}$) and, for each $\delta > 0$, let τ_δ be the first time t that $\eta'|_{[\tau,\infty)}$ hits $\partial B(w,\delta)$. Then the law of $\delta^{-1}(\eta'|_{[\tau,\tau_\delta]} - w)$ reparameterized by capacity converges weakly as $\delta \to 0$ to that of an $\mathrm{SLE}_6(2;2)$ process in $\mathbf{H}$ from 0 to ∞ with force points located at 0^- and 0^+ with respect to the topology of local uniform convergence.*

See Figure 5.4 for an illustration of the statement of Lemma 5.8. The proof of Lemma 5.8 will make use of the ideas developed in [MS12a, MS12b, MS12c, MS13a]. We will not give an in-depth introduction to imaginary geometry here, but rather refer the reader to the introductions of these articles as well as to [MW13, Section 2] for background. For $\kappa \in (0,4)$ and $\kappa' = 16/\kappa \in (4,8)$, we will make use of the following notation (which



Figure 5.4: Illustration of Lemma 5.8 and the setup of its proof. Shown is the time-reversal $\overline{\eta}'$ of an SLE_6 process η' in $\mathbf{H}$ from 0 to ∞ stopped at the first time $\overline{\tau}$ that it hits a given bounded interval $[x, y]$ in $\mathbf{R}$ containing a neighborhood of 0 on the event that $w = \overline{\eta}'(\overline{\tau}) \in [0, y]$. We show in Lemma 5.8 that if τ denotes the last time t that η' hits w and τ_δ is the first time t after time τ that η' hits $\partial B(w, \delta)$ then the law of $\delta^{-1}(\eta'(t) - w)$ for $t \in [\tau, \tau_\delta]$ converges as $\delta \to 0$ to an $\mathrm{SLE}_6(2; 2)$ process. The ρ values of $(2; 2)$ are special because they correspond to SLE_6 conditioned not to hit $\partial\mathbf{H}$ in the sense described in [MS12b]. Shown on the right is an initial segment of the right boundary η_R of $\overline{\eta}'$ as well as the left boundary $\widetilde{\eta}$ of $\overline{\eta}'|_{[0,\overline{\tau}]}$. In particular, $\widetilde{\eta}$ and η_R starting from the last time it hits $[0, y]$ together give the outer boundary of $\overline{\eta}'|_{[0,\overline{\tau}]}$. If we view $\overline{\eta}'$ as the counterflow line of a GFF h on $\mathbf{H}$ from ∞ to 0 with boundary conditions $\lambda' - \pi\chi$ (resp. $-\lambda' + \pi\chi$) on $\mathbf{R}_+$ (resp. $\mathbf{R}_-$), then η_R is the flow line of h starting from 0 of angle $-\pi/2$ and $\widetilde{\eta}$ is the flow line of the GFF given by restricting h to the left component of $\mathbf{H} \setminus \eta_R$ starting from the rightmost intersection point of η_R in $[x, y]$ with angle $\pi/2$.

matches that used in [MS12a, MS12b, MS12c, MS13a]):

$$\lambda = \frac{\pi}{\sqrt{\kappa}}, \quad \lambda' = \frac{\pi}{\sqrt{\kappa'}} = \lambda - \frac{\pi}{2}\chi, \quad \chi = \frac{2}{\sqrt{\kappa}} - \frac{\sqrt{\kappa}}{2}. \tag{5.3}$$

We recommend that the reader reference [MW13, Figure 2.5] while reading the proof of Lemma 5.8.

Proof of Lemma 5.8. We let $\kappa = 8/3$, $\kappa' = 16/\kappa = 6$, and will write λ, λ', and χ for the constants in (5.3) with these values of κ and κ'. By [MS12a, Theorem 1.1], we can view $\overline{\eta}'$ as the counterflow line from ∞ to 0 of a GFF h on $\mathbf{H}$ with boundary conditions given by $-\lambda' + \pi\chi = -\lambda + \frac{3}{2}\pi\chi$ (resp. $\lambda' - \pi\chi = \lambda - \frac{3}{2}\pi\chi$) on $\mathbf{R}_-$ (resp. $\mathbf{R}_+$). Moreover, by [MS12a, Theorem 1.4] we have that the left (resp. right) boundary of $\overline{\eta}'$ is given by the flow line η_L (resp. η_R) of h starting from 0 with angle $\pi/2$ (resp. $-\pi/2$). That is, η_L (resp. η_R) is the flow line of $h + \pi\chi/2$ (resp. $h - \pi\chi/2$) from 0 to ∞. By [MS12a, Theorem 1.1], the law of η_L (resp. η_R) is that of an $\mathrm{SLE}_\kappa(\kappa - 4; \kappa/2 - 2) = \mathrm{SLE}_{8/3}(-4/3; -2/3)$ (resp.

$\mathrm{SLE}_\kappa(\kappa/2-2;\kappa-4) = \mathrm{SLE}_{8/3}(-2/3;-4/3))$ process in $\mathbf{H}$ from 0 to ∞ with force points located at 0^- and 0^+. (See also [MW13, Figure 2.5].) In particular, η_L almost surely hits $\mathbf{R}_-$ but not $\mathbf{R}_+$ and likewise η_R almost surely hits $\mathbf{R}_+$ and not $\mathbf{R}_-$; recall that $\kappa/2-2$ is the critical ρ-value at or above which $\mathrm{SLE}_\kappa(\rho)$ does not hit the boundary.

Since $\overline{\eta}'$ visits the points on η_L (resp. η_R) in the reverse order in which they are visited by η_L (resp. η_R), we have that w is either equal to the last intersection of η_L with $[x,0]$ or to the last intersection of η_R with $[0,y]$. These two possibilities correspond to when $w \in [x,0]$ or $w \in [0,y]$. We shall assume without loss of generality that we are working on the event that the latter holds. Let $\widetilde{\eta}_L$ be the flow line with angle $\pi/2$ of the GFF given by restricting h to the component of $\mathbf{H} \setminus \eta_R$ which is to the left of η_R. It follows[4] from [DMS14, Proposition 6.7] that the two segments of η_R (which correspond to before and after the path hits w or, equivalently, before and after the path starts the excursion that it makes over y) translated by $-w$ and rescaled by the factor δ^{-1} converge as $\delta \to 0$ to a pair of flow lines η_1, η_2 of a GFF on $\mathbf{H}$ with boundary conditions $\lambda + \pi\chi/2 = \sqrt{2/3}\pi$ (resp. $-3\lambda + 5\pi\chi/2 = -\sqrt{2/3}\pi$) on $\mathbf{R}_+$ (resp. $\mathbf{R}_-$) with respective angles $\theta_1 = -\pi/2$, $\theta_2 = 2\lambda/\chi - \pi/2 = 5\pi/2$. Moreover, $\widetilde{\eta}_L$ translated by $-w$ and rescaled by δ^{-1} converges along with the two segments of η_R to the flow line of the same limiting GFF used to generate η_1, η_2 with angle $\pi/2$ (since the angle gap between $\widetilde{\eta}_L$ and η_R is equal to π, so the angle gap between $\widetilde{\eta}$ and η_1 is the same). In particular,

- The marginal law of η_1 is that of an $\mathrm{SLE}_\kappa(\kappa-2) = \mathrm{SLE}_{8/3}(2/3)$ process with a single force point located at 0^-.
- The marginal law of η_2 is that of an $\mathrm{SLE}_\kappa(\kappa-4;2) = \mathrm{SLE}_{8/3}(-4/3;2)$ process with force points located at 0^-, 0^+.
- The conditional law of η_2 given η_1 is that of an $\mathrm{SLE}_\kappa(\kappa-4) = \mathrm{SLE}_{8/3}(-4/3)$ process with force point located at 0^-.
- The conditional law of $\widetilde{\eta}$ given η_1, η_2 is that of an $\mathrm{SLE}_\kappa(\kappa/2-2;-\kappa/2) = \mathrm{SLE}_{8/3}(-2/3;-4/3)$ process with force points located immediately to the left and right of its starting point.

As $\kappa = 8/3$ and $\kappa' = 6$, we note that

$$\left(-3\lambda + \frac{5\pi\chi}{2}\right) - \pi\chi = -3\lambda + \frac{3\pi\chi}{2} = -3\lambda' \quad \text{and}$$
$$\left(\lambda + \frac{\pi\chi}{2}\right) + \pi\chi = \lambda + \frac{3\pi\chi}{2} = 3\lambda'.$$

[4] [DMS14, Proposition 6.7] is stated in the setting of $\mathrm{SLE}_\kappa(\rho)$ processes with a single boundary force point; however, the same proof goes through verbatim to describe the local behavior of the start point an excursion which straddles a given boundary point for $\mathrm{SLE}_\kappa(\rho_1;\rho_2)$ processes.

Therefore by [MS12a, Theorem 1.4], we have that $\widetilde{\eta}$ and η_1 respectively give the left and right boundaries of the counterflow line from ∞ to 0 which, by the form of the boundary data, is an $\mathrm{SLE}_6(2;2)$ process with force points located immediately to the left and right of ∞. The result thus follows by the reversibility of $\mathrm{SLE}_6(2;2)$ established in [MS12c]. □

Lemma 5.9. *Suppose that η' is a whole-plane* SLE_6 *process from 0 to ∞ in $\mathbf{C}$. Let τ be a stopping time for η' so that $\eta'([0,\tau])$ is almost surely bounded and let $\overline{\sigma}$ be a stopping time for the time-reversal $\overline{\eta}'$ of η' given $\eta'|_{[0,\tau]}$ such that $\overline{\eta}'|_{[0,\overline{\sigma}]}$ does not hit $\eta'([0,\tau])$ almost surely. Let $\overline{\tau}$ be the first time that $\overline{\eta}'$ hits $\eta'([0,\tau])$. Let ψ be the unique conformal map from the unbounded component of $\mathbf{C}\setminus\eta'([0,\tau])$ to $\mathbf{H}$ which sends ∞ to i and $\eta'(\tau)$ to 0. Then law of the time-reversal of $\delta^{-1}\psi(\overline{\eta}'|_{[0,\overline{\tau}]})$ parameterized by capacity converges weakly respect to the topology of local uniform convergence as $\delta\to 0$ to that of a chordal* $\mathrm{SLE}_6(2;2)$ *process from 0 to ∞ with force points located at 0^- and 0^+.*

Proof. This follows by combining the locality property of whole-plane SLE_6 with Lemma 5.8. □

Lemma 5.10. *Suppose that η' is a whole-plane* SLE_6 *process from 0 to ∞ in $\mathbf{C}$. Let τ be a stopping time for η' such that $\eta'([0,\tau])$ is bounded almost surely. Let $\overline{\eta}'$ be the time-reversal of η' and let $\overline{\tau}$ be the first time that $\overline{\eta}'$ hits $\eta'([0,\tau])$. Then $\overline{\eta}'(\overline{\tau})$ is distributed according to harmonic measure as seen from ∞ on the boundary of the unbounded component of $\mathbf{C}\setminus\eta'([0,\tau])$.*

Proof. This is a consequence of the locality for whole-plane SLE_6, the domain Markov property, and conformal invariance. □

Proof of Proposition 5.7. We will first prove a version of the proposition in which we have not conditioned on E_ϵ or $\mathcal{F}$. That is, we will first argue that as $\epsilon\to 0$ (but $u>0$ fixed), we have that the joint law of $(\mathbf{H},\widetilde{h})$ and $\widetilde{\eta}'$ under $\mathbf{P}_u$ converges to a pair consisting of a $\sqrt{8/3}$-quantum wedge normalized to assign quantum boundary length 1 to $[-1,0]$ and an independent chordal SLE_6 process in $\mathbf{H}$ from 0 to ∞.

We will establish this using Lemma 5.6. Suppose that $\widehat{\mathcal{W}}=(\mathbf{H},\widehat{h},0,\infty)$ is a $\sqrt{8/3}$-quantum wedge and that $\widehat{\eta}'$ is a chordal SLE_6 process from 0 to ∞ sampled independently of $\widehat{h}$. Let $(\widehat{f}_t)$ denote the forward centered Loewner flow associated with $\widehat{\eta}'$ and, for each s, we let $\widehat{X}_s$ denote the change in the boundary length of $h\circ\widehat{f}_s^{-1}+Q\log|(\widehat{f}_s^{-1})'|$ relative to $s=0$. Assume that $\widehat{\eta}'$ is parameterized according to quantum natural time. By [DMS14, Corollary 1.20], we have that $\widehat{X}_s$ evolves as a 3/2-stable Lévy process with only downward jumps. Moreover, by [DMS14, Theorem 1.19], we have that conditional on the realization of the entire process $\widehat{X}_s$, the components of $\mathbf{H}\setminus\widehat{\eta}'$ (viewed as quantum surfaces) are conditionally independent quantum disks given their boundary length and the boundary length of each such disk is given by the corresponding jump of $\widehat{X}_s$.

We can construct a coupling of the surface near $\eta'(\mathfrak{t})$ and $\widehat{\mathcal{W}}$ as follows. Let X_s denote the quantum length of the outer boundary of $\eta'([0,\mathfrak{t}+s])$ relative to the quantum length of the outer boundary of $\eta'([0,\mathfrak{t}])$. In other words, we normalize so that $X_0 = 0$. Fix $\delta > 0$. Lemma 5.6 implies that there exists $s_0 > 0$ such that we can find a coupling of X and $\widehat{X}$ so that the event $A = \{X|_{[0,s_0]} = \widehat{X}|_{[0,s_0]}\}$ satisfies $\mathbf{P}_u[A] \geq 1-\delta$. Note that the conditional law of the quantum disks cut off by η' in the time interval $[0,s_0]$ given $X|_{[0,s_0]}$ is the same as that for the quantum disks cut off by $\widehat{\eta}'$ in the time interval $[0,s_0]$ given $\widehat{X}|_{[0,s_0]}$. Consequently, we can couple together these quantum disks so that they are the same (equal as boundary-marked quantum surfaces and with the same orientation) on A. We sample the remainder of the quantum disks conditionally independently. On A^c, we sample all of the quantum disks conditionally independently given $X|_{[0,s_0]}$ and $\widehat{X}|_{[0,s_0]}$.

Let $\widetilde{K}$ (resp. $\widehat{K}$) be the region cut off from ∞ by $\widetilde{\eta}'([0,s_0])$ (resp. $\widehat{\eta}'([0,s_0])$). On A, we have that $(\widetilde{K},\widetilde{h},0)$ and $(\widehat{K},\widehat{h},0)$ are equal as (marked) quantum surfaces and $(\mathbf{H}\setminus\widetilde{K},\widetilde{h})$ and $(\mathbf{H}\setminus\widehat{K},\widehat{h})$ are conditionally independent as quantum surfaces given $X_{s_0} = \widehat{X}_{s_0}$. Let $\widetilde{V}$ (resp. $\widehat{V}$) be the component of the interior of $\widetilde{K}$ (resp. $\widehat{K}$) which contains 0. Then there exists a conformal map $\widetilde{\psi}\colon \widetilde{V} \to \widehat{V}$ which takes 0 to 0 since $(\widetilde{V},\widetilde{h},0)$ and $(\widehat{V},\widehat{h},0)$ are equal as (marked) quantum surfaces. By scaling, we can take the embedding of $(\mathbf{H},\widehat{h})$ so that $\widetilde{\psi}'(0) = 1$. It therefore follows that $|\widetilde{\psi}(z) - z| \to 0$ as $z \to 0$, hence it is easy to see that the total variation distance between the laws of the restrictions of the two surfaces to $\mathbf{H} \cap B(0,\delta)$ converges to 0 as $\delta \to 0$.

We have now proved the part of the proposition which involves the behavior of the surface near $\eta'(\mathfrak{t})$. The same proof works if we condition on E_ϵ and $\mathcal{F}$ since this conditioning does not change the local behavior of the surface near 0. We shall now assume that we are conditioning on both E_ϵ and $\mathcal{F}$. To finish the proof, we need to describe the behavior of $\widetilde{\eta}'$ near $\widetilde{\eta}'(\mathfrak{t})$. Let $z_0 = \varphi(\infty)$ and let $\overline{\tau}$ be the first time that $\overline{\eta}'$ hits $\eta'([0,\mathfrak{t}])$. Lemma 5.10 then implies that the distribution of $\varphi(\overline{\eta}'(\overline{\tau}))$ is equal to the distribution on $[-1,0]$ induced by harmonic measure as seen from z_0 (normalized to be a probability). Since we know that $z_0 \to \infty$ as $\epsilon \to 0$, it follows that the distribution of $\varphi(\overline{\eta}'(\overline{\tau}))$ converges to the uniform distribution as $\epsilon \to 0$. Moreover, that the law of the path is as claimed follows from Lemma 5.9. □

Lemma 5.11. *There exists $p_0 > 0$ such that for each $u > 0$ we have that*

$$\mathbf{P}_u[F_\epsilon \,|\, E_\epsilon] \to p_0 \quad \text{as} \quad \epsilon \to 0.$$

Proof. This is obvious from the representation of the conditional law of the configuration near the tip described in Proposition 5.7. □

We are now ready to combine everything to complete the proof of Theorem 5.1.

Proof of Theorem 5.1. As mentioned above, we will complete the proof of the theorem by establishing (5.1) and (5.2). We now introduce the topology respect to which we will establish the weak limits in (5.1) and (5.2). (As mentioned earlier, the exact topology is not important.)

We will first define a metric on the space $\mathbf{X}$ of quantum surfaces which are homeomorphic to $\mathbf{D}$. We can associate with each quantum surface $\mathcal{S} = (D, h)$ a probability measure on random distributions $\widetilde{h}$ by picking $z \in D$ according to the quantum measure, $\theta \in [0, 2\pi]$ uniformly at random and independently of z, letting $\varphi \colon D \to \mathbf{D}$ be the unique conformal transformation with $\varphi(z) = 0$ and $\varphi'(0)/|\varphi'(0)| = e^{i\theta}$, and then taking $\widetilde{h} = h \circ \varphi^{-1} + Q \log |(\varphi^{-1})'|$. Clearly, if $\mathcal{S}_1 = (D_1, h_1)$ and $\mathcal{S}_2 = (D_2, h_2)$ are equivalent as quantum surfaces then the corresponding random distributions $\widetilde{h}_1$ and $\widetilde{h}_2$ on $\mathbf{D}$ have the same law. We define our metric d on $\mathbf{X}$ by taking $d(\mathcal{S}_1, \mathcal{S}_2)$ for $\mathcal{S}_1, \mathcal{S}_2 \in \mathbf{X}$ to be given by the sum over $j \in \mathbf{N}$ of 2^{-j} times the Prokhorov distance between the laws of the restrictions of $\widetilde{h}_1$ and $\widetilde{h}_2$ to $B(0, 1 - 2^{-j})$. We can also extend the definition of $\mathbf{X}$ to the setting of k-marked quantum surfaces $\mathbf{X}_k$ which are homeomorphic to $\mathbf{D}$ by taking the sum over $j \in \mathbf{N}$ of 2^{-j} times the Prokhorov distance between the joint law of the restriction of $\widetilde{h}_1$ to $B(0, 1 - 2^{-j})$ and its marked points and the joint law of the restriction of $\widetilde{h}_2$ to $B(0, 1 - 2^{-j})$ and its marked points.

We next recall from [DMS14, MS15b] (as well as Section 4) that we can encode an SLE_6 trajectory on top of a $\sqrt{8/3}$-LQG sphere using the time-reversal e of a 3/2-stable Lévy excursion and the ordered collection of marked, oriented quantum disks that the path cuts off from its target point. Therefore we can view the path-decorated, beaded quantum surface which corresponds to the region separated by an SLE_6 from its target point as well as the path stopped at a given time as a random variable which takes values in $\mathbf{X}^*$ which is the product of the subspace of càdlàg functions in ℓ^∞ with only downward jumps and $(\mathbf{X}_1 \times \{0, 1\})^{\mathbf{N}}$; we equip $\mathbf{X}^*$ with the product topology. Indeed, the time-reversed Lévy excursion naturally is an element of the former space and the ordered collection of marked, oriented quantum disks naturally take values in the latter space (the orientation corresponds to the extra bit).

Fix $\epsilon > 0$. For any interval $I = (x_1, x_2)$ with $0 < x_1 < x_2$ and any interval $J = (t_1, t_2)$ with $0 < t_1 < t_2$ we have that $\mathsf{M}^2_{\mathrm{SPH,W}}[X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] \in (0, \infty)$ so that $\mathsf{M}^2_{\mathrm{SPH,W}}$ conditioned on $X_{\mathfrak{t}} \in I$ and $\mathfrak{t} \in J$ makes sense as a probability measure. We can thus write

$$\mathsf{M}^2_{\mathrm{SPH,W}}[E_\epsilon, X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] = \mathsf{M}^2_{\mathrm{SPH,W}}[E_\epsilon \,|\, X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] \mathsf{M}^2_{\mathrm{SPH,W}}[X_{\mathfrak{t}} \in I, \mathfrak{t} \in J].$$

Lemma 5.4 implies that for all $\epsilon \in (0, x_1)$ we have that

$$\frac{\epsilon}{x_2} \leq \mathsf{M}^2_{\mathrm{SPH,W}}[E_\epsilon \,|\, X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] \leq \frac{\epsilon}{x_1}. \tag{5.4}$$

We let

$$G_j = \{2^{j-1} \leq \mathfrak{t} \leq 2^j\} \quad \text{for each} \quad j \in \mathbf{Z}.$$

Suppose that $f\colon \mathbf{X}^* \times \mathbf{X}^* \times \mathbf{X} \to \mathbf{R}$ is a bounded, continuous function. Let $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ be as in the statement of Theorem 5.1. Lemma 5.5 implies that the $\mathsf{M}^2_{\mathrm{SPH,W}}$ conditional laws of $\mathcal{X}$, $\overline{\mathcal{X}}$, and $\mathcal{B}$ are the same if we condition on $X_{\mathfrak{t}}$ or if we condition on both $X_{\mathfrak{t}}$ and E_ϵ. It therefore follows from (5.4) that for each $j \in \mathbf{Z}$ we have

$$\epsilon^{-1} \int f(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B}) \mathbf{1}_{E_\epsilon \cap G_j} d\mathsf{M}^2_{\mathrm{SPH,W}} \to \int f(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B}) \mathbf{1}_{G_j} d\mathsf{M}^2_{\mathrm{SPH,D}} \quad \text{as} \quad \epsilon \to 0. \tag{5.5}$$

Let $\mathcal{Y}$, $\overline{\mathcal{Y}}$, and $\mathcal{Q}$ be as in the statement of Proposition 5.3 and let $\mathcal{Q}^*$ be the component of $\mathcal{Q}$ with the largest quantum area (there is almost surely a unique such component). Then Proposition 5.3 implies that $\mathsf{M}^2_{\mathrm{SPH,W}}$ is invariant under the operation of swapping $\mathcal{Y}$ and $\overline{\mathcal{Y}}$. Since the event F_ϵ is also defined in a way which is symmetric under swapping $\mathcal{Y}$ and $\overline{\mathcal{Y}}$, we have that the measure $\mathbf{1}_{F_\epsilon} d\mathsf{M}^2_{\mathrm{SPH,W}}$ is symmetric in the same sense. Thus to finish the proof it suffices to show that there exists a constant $c_0 > 0$ such that for any f as above,

$$\epsilon^{-1} \int f(\mathcal{Y}, \overline{\mathcal{Y}}, \mathcal{Q}^*) \mathbf{1}_{F_\epsilon \cap G_j} d\mathsf{M}^2_{\mathrm{SPH,W}} \to c_0 \int f(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B}) \mathbf{1}_{G_j} d\mathsf{M}^2_{\mathrm{SPH,D}} \quad \text{as} \quad \epsilon \to 0. \tag{5.6}$$

Note that we in fact have that $\mathcal{X} = \mathcal{Y}$. Moreover, it follows from Proposition 5.7 that for each $\delta > 0$ and $j \in \mathbf{Z}$ we have that

$$\mathsf{M}^2_{\mathrm{SPH}}[d(\overline{\mathcal{X}}, \overline{\mathcal{Y}}) \geq \delta \,|\, E_\epsilon, G_j] \to 0 \quad \text{as} \quad \epsilon \to 0.$$

Lemma 5.11 implies that the same is true when we condition on F_ϵ instead of E_ϵ. Therefore for f bounded and continuous as above, we have that

$$\epsilon^{-1} \int |f(\mathcal{Y}, \overline{\mathcal{Y}}, \mathcal{Q}^*) - f(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})| \mathbf{1}_{F_\epsilon \cap G_j} d\mathsf{M}^2_{\mathrm{SPH,W}} \to 0 \quad \text{as} \quad \epsilon \to 0$$

for each $j \in \mathbf{Z}$. Therefore it suffices to prove (5.6) with $\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B}$ in place of $\mathcal{Y}, \overline{\mathcal{Y}}, \mathcal{Q}^*$. Assume that I, J are as above. Then we have that

$$\begin{aligned} &\mathsf{M}^2_{\mathrm{SPH,W}}[F_\epsilon, X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] \\ =&\mathsf{M}^2_{\mathrm{SPH,W}}[F_\epsilon \,|\, E_\epsilon, X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] \mathsf{M}^2_{\mathrm{SPH,W}}[E_\epsilon \,|\, X_{\mathfrak{t}} \in I, \mathfrak{t} \in J] \mathsf{M}^2_{\mathrm{SPH,W}}[X_{\mathfrak{t}} \in I, \mathfrak{t} \in J]. \end{aligned}$$

By Lemma 5.11, we have that the first term on the right hand side converges to p_0 as $\epsilon \to 0$. Therefore (5.6) follows from (5.5). □

As a consequence of the analysis which gave the proof of Theorem 5.1, we will also be able to identify the conditional law of $\mathcal{B}$ under $\mathsf{M}^2_{\mathrm{SPH,D}}$ given its boundary length. This will be rather important for us later in this article because, as we will explain now, it implies that $\partial\mathcal{B}$ is almost surely conformally removable.

Proposition 5.12. *Suppose that $\mathcal{B}$ is as in Theorem 5.1.*

(i) Conditionally on its boundary length, the law of $\mathcal{B}$ under $\mathsf{M}^2_{\mathrm{SPH,D}}$ is that of a quantum disk.

(ii) Suppose that $(\mathbf{C}, h)$ is any embedding of $(\mathcal{S}, x, y)$ distributed according to $\mathsf{M}^2_{\mathrm{SPH,D}}$. Then the image of the embedding of $\partial\mathcal{B}$ is almost surely conformally removable.

Proof. First of all, we know from Lemma 5.5 that the conditional law of $\mathcal{B}$ given $\mathfrak{t}$ and $X_{\mathfrak{t}}$ is the same as the conditional law of $\mathcal{B}$ given $\mathfrak{t}$, $X_{\mathfrak{t}}$, and E_ϵ. We define $d(\mathcal{S}^1, \mathcal{S}^2)$ as in the proof of Theorem 5.1. The conditional law of the segment $\widetilde{\eta}'$ of η' which is contained in $\mathcal{B}$ and connects $\eta'(\mathfrak{t})$ to $\overline{\eta}'(\overline{\mathfrak{t}})$ (i.e., the segment indicated in green in Figure 5.2) is given by a chordal SLE_6 process independent of $\mathcal{B}$. Let $\widetilde{\mathcal{B}}$ be the component of $\mathcal{B} \setminus \widetilde{\eta}'$ with the longest quantum boundary length. Then we know that $\widetilde{\mathcal{B}}$ is a quantum disk conditional on its boundary length because it is a complementary component of η' on $\mathcal{S}$. The result follows because the d-distance between $\widetilde{\mathcal{B}}$ and $\mathcal{B}$ tends to 0 in probability as $\epsilon \to 0$ while, as we mentioned above, the law of $\mathcal{B}$ does not change with ϵ. This proves that the conditional law of $\mathcal{B}$ given its boundary length is a quantum disk.

Assume that $\mathcal{B}$ is parameterized by $(\mathcal{S}, \widetilde{h})$. Then we know that the law of $\widetilde{h}$ is absolutely continuous with respect to a free boundary GFF on $[a, b] \times [0, 2\pi]$ for any $a, b \in \mathbf{R}$ with $a < b$. Likewise, we know that the law of h in any bounded region $U \subseteq \mathbf{C}$ is mutually absolutely continuous with respect to the law of the restriction to U of a free boundary GFF on a bounded domain with $\overline{U} \subseteq V$. Therefore it follows from [MS13b, Theorem 8.1] that the map $\varphi\colon [a, b] \times [0, 2\pi] \to \mathbf{C}$ which corresponds to the embedding of $\mathcal{B}$ into $\mathbf{C}$ is almost surely Hölder continuous in $[a, b] \times [0, 2\pi]$. By [DMS14, Corollary 5.8], we know that the points on $\partial\mathcal{B}$ which correspond to $\pm\infty \in \partial\mathcal{S}$ are uniformly distributed according to the quantum boundary measure on $\partial\mathcal{B}$. By resampling them, it follows that the embedding of $\mathcal{B}$ into $\mathbf{C}$ is almost surely a Hölder domain. Therefore it follows from [JS00, Corollary 2] that the embedding of $\partial\mathcal{B}$ into $\mathbf{C}$ is almost surely conformally removable. □

6 Quantum natural time $\mathrm{QLE}(8/3, 0)$

The purpose of this section is to construct a "quantum natural time" version of $\mathrm{QLE}(8/3, 0)$. This is a variant of the process constructed in [MS13b] where the approximations involve resampling the tip of the SLE_6 at δ-units of quantum natural time as opposed to δ-units of capacity time. The construction that we will give here describes a growth process on a quantum sphere, which is also in contrast to [MS13b] in which the process is constructed on an (unscaled) quantum cone. The present construction also generalizes to the setting of quantum cones, but we will focus on the sphere case. Throughout this article, whenever we refer to the process $\mathrm{QLE}(8/3, 0)$ we mean the one constructed just below unless explicitly stated otherwise. As in the case of the construction given in [MS13b], we will begin in Section 6.1 by introducing the

approximations to QLE$(8/3, 0)$. We will then show in Section 6.2 that the subsequential limits of these approximations have the following properties which will be important later on:

- The bubbles swallowed by the process have the same Poissonian structure as the bubbles swallowed by an SLE_6,
- The evolution of the quantum boundary length of the complementary component which contains the target point is the same as in the case of an SLE_6 (the time-reversal of a 3/2-stable Lévy excursion with only positive jumps), and
- The law of the region which contains the target point is the same as in the case of SLE_6 (that of a quantum disk weighted by its area).

6.1 Approximations to QLE$(8/3, 0)$

Fix $\delta > 0$ and suppose that $e \colon [0, T] \to \mathbf{R}_+$ is a sample picked from the excursion measure N for a 3/2-stable Lévy process with only positive jumps as described in Section 4.1. We define the δ-approximation to QLE$(8/3, 0)$ associated with the excursion e as follows. First, we let $(\mathcal{S}, x, y)$ be a doubly-marked quantum sphere constructed from e and let $\eta'_1 = \eta'$ be the associated whole-plane SLE_6 from x to y, as described in Section 4.1, with the quantum natural time parameterization. For concreteness, we will take the embedding so that $(\mathcal{S}, x, y) = (\mathbf{C}, h, 0, \infty)$ where 0 (resp. ∞) corresponds to x (resp. 0) and the scaling factor is determined so that the quantum mass of $\mathbf{D}$ is equal to 1/2 the total quantum mass of $\mathcal{S}$.

We define a growth process Γ^δ inductively as follows. First, we take Γ^δ_t to be the complement of the unbounded component of $\mathbf{C} \setminus \eta'([0, t])$ for each $t \in [0, \delta]$. We also let $g^\delta_t \colon \mathbf{C} \setminus \Gamma^\delta_t \to \mathbf{C} \setminus \mathbf{D}$ be the unique conformal map with $|g^\delta_t(z) - z| \to 0$ as $z \to \infty$. Fix $j \in \mathbf{N}$ and suppose that we have defined paths $\eta'_1, \ldots, \eta'_j$ and a growing family of hulls Γ^δ with associated uniformizing conformal maps (g^δ_t) for $t \in [0, j\delta]$ such that the following hold:

- The conditional law of the surface parameterized by the complement of $\Gamma^\delta_{j\delta}$ given its quantum boundary length is the same as in the setting of ordinary SLE_6. That is, it is given by a quantum disk with the given boundary length weighted by its area.
- $\eta'_j(j\delta)$ is distributed uniformly according to the quantum boundary measure on $\partial\Gamma^\delta_{j\delta}$ conditional on $\Gamma_{j\delta}$ (as a path decorated surface).
- The law of the components separated from the target point by time $j\delta$ is the same as in the case of whole-plane SLE_6. That is, they are given by conditionally independent quantum disks given their boundary lengths.

We then let η'_{j+1} be an independent radial SLE_6 starting from a point on $\partial\Gamma^\delta_{j\delta}$ which is chosen uniformly from the quantum boundary measure conditionally independently of everything else (i.e., we resample the location of the tip $\eta'(j\delta)$). For each $t \in [j\delta, (j+1)\delta]$, we also let Γ^δ_t be the complement of the unbounded component of $\mathbf{C} \setminus (\Gamma^\delta_{j\delta} \cup \eta'_{j+1}([0,t]))$. Then by the construction, all three properties described above are satisfied by the process up to time $(j+1)\delta$.

By resampling the surface parameterized by the complement of $\Gamma^\delta_{j\delta}$ at each stage, we can arrange so that the quantum boundary length of this component at each time t is given by the time-reversal $e(T-t)$ of e. In particular, the jumps of e then correspond to the quantum boundary length of the quantum disks swallowed by Γ^δ.

It will be convenient to encode Γ^δ in terms of a radial Loewner flow. That is, if for each $t \geq 0$ we let $s(t)$ be the quantum natural time elapsed by Γ^δ at the first time that the capacity as seen from ∞ reaches t then there exists a measure ν_δ on $\partial\mathbf{D} \times [0,\infty)$ whose second-coordinate marginal is given by Lebesgue measure such that

$$g^\delta_{s(t)}(z) = z + \iint_{\partial\mathbf{D}\times[0,t]} g^\delta_{s(u)}(z) \frac{w + g^\delta_{s(u)}(z)}{w - g^\delta_{s(u)}(z)} d\nu_\delta(w,u) \quad \text{for all} \quad t \geq 0.$$

Since the growth during each of the δ-length time intervals is given by a segment of an independent radial SLE_6 process, it follows that there exists a standard Brownian motion B and a sequence of real numbers (ξ_k) such that with $W = \sqrt{6}B$ we have that

$$d\nu_\delta(\theta,t) = \sum_k \mathbf{1}_{[k\delta,(k+1)\delta]}(s(t)) \delta_{e^{i(W_t+\xi_k)}} dt$$

where δ_x denotes the Dirac mass supported at $x \in \partial\mathbf{D}$ and dt denotes Lebesgue measure on $\mathbf{R}_+$.

We emphasize that the δ-approximation to $\mathrm{QLE}(8/3,0)$ satisfies the following:

- The bubbles which it separates from y are conditionally independent quantum disks given their boundary lengths,
- The boundary length of the complementary component which contains y at time t is equal to the time-reversal of e at time t and is conditionally independent of everything else given its boundary length, and
- The conditional law of the region which contains y given its quantum boundary length is the same as in the case of SLE_6. That is, it is given by a quantum disk with the given boundary length weighted by its area.

6.2 Subsequential limits

We are now going to construct subsequential limits of the δ-approximations to QLE$(8/3, 0)$ described just above. We will pick the subsequence so that the limit we obtain satisfies the three properties listed just above.

For each $T > 0$, we let ν_δ^T be the restriction of ν_δ to $\partial\mathbf{D} \times [0, T]$. Since $\partial\mathbf{D} \times [0, T]$ is compact and each ν_δ^T has total mass equal to T, it follows that the law on random measures (ν_δ^T) is tight in $\delta > 0$ (but T fixed) with respect to the weak topology on measures on $\partial\mathbf{D} \times [0, T]$. For each $\delta > 0$, we let $(\mathcal{S}_\delta, x_\delta, y_\delta)$ be the doubly-marked quantum sphere associated with ν_δ. We take the embedding of $(\mathcal{S}_\delta, x_\delta, y_\delta)$ to be given by $(\mathbf{C}, h_\delta, 0, \infty)$, as in Section 6.1, where the scaling factor is chosen so that the quantum mass of $\mathbf{D}$ is equal to $1/2$ the quantum mass of $\mathcal{S}$.

For each $0 < a_1 < a_2 < \infty$, we let E_{a_1,a_2}^δ be the event that the area of $\mathcal{S}_\delta$ is contained in $[a_1, a_2]$. We note that E_{a_1,a_2}^δ is an event of positive and finite $\mathsf{M}_{\mathrm{SPH}}^2$ measure so that the conditional law of $(\mathbf{C}, h_\delta, 0, \infty)$ given E_{a_1,a_2}^δ makes sense as a probability measure. Since the law of h_δ conditioned on E_{a_1,a_2}^δ does not depend on δ, it follows that there exists a sequence (δ_k) with $\delta_k > 0$ for all k and $\delta_k \to 0$ as $k \to \infty$ such that the joint law of h_δ and $\nu_{\delta_k}^T$ given $E_{a_1,a_2}^{\delta_k}$ converges weakly to a limiting law as $k \to \infty$. (We take the notion of convergence of h_δ to be that its coordinates converge when expanded in terms of an orthonormal basis of $H(\mathbf{C})$ consisting of $C_0^\infty(\mathbf{C})$ functions.) The Skorohod embedding theorem implies that we can find a coupling of the laws of h_{δ_k} and $\nu_{\delta_k}^T$ given $E_{a_1,a_2}^{\delta_k}$ (as k varies) with a doubly-marked quantum sphere $(\mathbf{C}, h, 0, \infty)$ conditioned on the event E_{a_1,a_2} that its area is contained in $[a_1, a_2]$ together with a measure ν^T on $\partial\mathbf{D} \times [0, T]$ whose second coordinate marginal is given by Lebesgue measure such that $h_{\delta_k} \to h$ as described above and $\nu_{\delta_k}^T \to \nu^T$ weakly almost surely as $k \to \infty$. By passing to a further (diagonal) subsequence, we can find a coupling of the laws of the $(\mathbf{C}, h_{\delta_k}, 0, \infty)$ (unconditioned) and (ν_{δ_k}) and $(\mathbf{C}, h, 0, \infty)$ and a measure ν on $\partial\mathbf{D} \times [0, \infty)$ whose second coordinate marginal is given by Lebesgue measure so that with ν^T equal to the restriction of ν to $\partial\mathbf{D} \times [0, T]$ for each $T \geq 0$ we have that $h_{\delta_k} \to h$ and $\nu_{\delta_k}^T \to \nu^T$ weakly almost everywhere for all $T > 0$. By [MS13b, Theorem 1.1], it follows that the radial Loewner flows driven by the ν_{δ_k} converge locally uniformly in time and space to the radial Loewner flow driven by ν as $k \to \infty$ as well.

Definition 6.1. Suppose that $(\mathcal{S}, x, y) = (\mathbf{C}, h, 0, \infty)$ is a doubly-marked quantum surface and that ν is a measure on $\partial\mathbf{D} \times [0, \infty)$ so that the joint law of $(\mathcal{S}, x, y)$ and ν is equal to any one of the subsequential limits constructed above. Then we refer to the increasing family Γ (modulo time parameterization) as QLE$(8/3, 0)$ where Γ is the growth process which corresponds to the radial Loewner flow driven by ν.

As explained in the introduction, it will be a consequence of the results of this work and [MS16a] that it is not necessary to pass along a subsequence (δ_k) in the construction of QLE$(8/3, 0)$.

Remark 6.2. As we will explain below, there are several natural parameterizations of time for QLE$(8/3, 0)$. We emphasize that the definition of QLE$(8/3, 0)$ given in Definition 6.1 does not carry with it a time parameterization. We in particular do not build the capacity parameterization into the definition because it depends on the embedding of the ambient surface into $\mathbf{C}$. Later, we will want to be able to cut a QLE$(8/3, 0)$ out of a surface and glue it into another and such an operation in general does not preserve capacity because it can lead to a different embedding. The time-parameterizations that we introduce shortly are, however, intrinsic to QLE$(8/3, 0)$ viewed as a growing family of quantum surfaces.

Suppose that Γ is a QLE$(8/3, 0)$ on a doubly-marked sphere $(\mathcal{S}, x, y)$. We say that a region U is swallowed by Γ if U is equal to the interior of $\Gamma_t \setminus \Gamma_{t-}$ for some t. The following proposition will imply that we can make sense of the quantum natural time parameterization of Γ.

Proposition 6.3. *Suppose that $(\mathcal{S}, x, y)$ is a doubly-marked quantum sphere with distribution $\mathsf{M}^2_{\mathrm{SPH}}$ and that Γ is a* QLE$(8/3, 0)$ *on $\mathcal{S}$ from x targeted at y. Then the joint law of the regions swallowed by Γ, ordered by the time at which they are swallowed and viewed as quantum surfaces, is the same as for a whole-plane* SLE_6 *on $\mathcal{S}$ connecting x to y.*

Proof. We shall assume that we are in the setting described at the beginning of this subsection. In particular, we take the embedding of $(\mathcal{S}, x, y)$ to be given by $(\mathbf{C}, h, 0, \infty)$.

For each $k \in \mathbf{N}$, we let Γ^{δ_k} be the growth process associated with ν_{δ_k}. We take Γ^{δ_k} to be parameterized by capacity as seen from ∞. By the construction of the δ_k-approximation to QLE$(8/3, 0)$, we know the law of the regions swallowed by Γ^{δ_k} ordered by t and viewed as quantum surfaces is the same as for a whole-plane SLE_6 on $(\mathcal{S}, x, y)$ connecting x to y. Our aim is to show that this result holds in the limit as $k \to \infty$.

For each k, we let $\mathcal{U}^{\delta_k}$ be the collection of surfaces swallowed by Γ^{δ_k} ordered by the time at which they are swallowed. We note that for each $\epsilon > 0$, the collection $\mathcal{U}^{\delta_k}_\epsilon$ which consists of those elements of $\mathcal{U}^{\delta_k}$ with quantum boundary length at least ϵ is finite almost surely. Indeed, this follows because for N almost every e we have that the number of jumps made by e of size at least ϵ is finite. Since the law of each of the $\mathcal{U}^{\delta_k}$ is the same for each k, by possibly passing to a further subsequence (and recoupling the laws using the Skorohod embedding theorem so that we have almost sure convergence) we have that each $|\mathcal{U}^{\delta_k}_\epsilon|$ converges almost surely to a finite limit as $k \to \infty$.

For each j we let $U^{\delta_k}_{j,\epsilon}$ be the jth element of $\mathcal{U}^{\delta_k}_\epsilon$, $z^{\delta_k}_{j,\epsilon}$ be a point chosen uniformly at random from the quantum measure restricted to $U^{\delta_k}_{j,\epsilon}$, and let $\varphi^{\delta_k}_{j,\epsilon} \colon \mathbf{D} \to U^{\delta_k}_{j,\epsilon}$ be the unique conformal map with $\varphi^{\delta_k}_{j,\epsilon}(0) = z^{\delta_k}_{j,\epsilon}$ and $(\varphi^{\delta_k}_{j,\epsilon})'(0) > 0$. (If $j > |\mathcal{U}^{\delta_k}_\epsilon|$ then we take $\varphi^{\delta_k}_{j,\epsilon} \equiv 0$.) Fix an orthonormal basis (ϕ_n) of $H(\mathbf{D})$ consisting of $C^\infty_0(\mathbf{D})$ functions. Each element of $\mathcal{U}^{\delta_k}_\epsilon$ is a quantum disk hence can be described by a distribution on $\mathbf{D}$ given

by $h_{\delta_k} \circ \varphi_{j,\epsilon}^{\delta_k} + Q \log |(\varphi_{j,\epsilon}^{\delta_k})'|$. We can express this distribution in terms of coordinates with respect to the orthonormal basis (ϕ_n). Therefore by possibly passing to a further, diagonal subsequence (and recoupling the laws using the Skorohod embedding theorem so that we have almost sure convergence) we have for each fixed j that the jth element of $\mathcal{U}_\epsilon^{\delta_k}$ almost surely converges to a limiting distribution on $\mathbf{D}$ in the sense that each of its coordinates with respect to (ϕ_n) converge almost surely (if j is larger than the number of elements of $\mathcal{U}_\epsilon^{\delta_k}$ then we take the jth element to be the zero distribution on $\mathbf{D}$). Combining, we have that $\mathcal{U}_\epsilon^{\delta_k}$ converges almost surely to a limit $\mathcal{U}_\epsilon$ which has the same law as each of the $\mathcal{U}_\epsilon^{\delta_k}$ in the sense described just above (number of elements converges and each element converges weakly almost surely).

We note that the laws of the $(\varphi_{j,\epsilon}^{\delta_k})'(0)$ are tight as $k \to \infty$. Indeed, if there was a uniformly positive chance that one of the $(\varphi_{j,\epsilon}^{\delta_k})'(0)$ is arbitrarily large for large k, then we would have that there is a uniformly positive chance that $(\mathcal{S}^k, x^k, y^k) = (\mathbf{C}, h_{\delta_k}, 0, \infty)$ assigns a uniformly positive amount of mass to $\mathbf{C} \setminus B(0, R)$ for each $R > 0$ and k large enough. This, in turn, would lead to the contradiction that the limiting surface $(\mathcal{S}, x, y)$ would have an atom at y with positive probability. It therefore follows that by passing to a further subsequence if necessary (and recoupling the laws using the Skorohod embedding theorem so that we have almost sure convergence), we can arrange so that each of the conformal maps $\varphi_{j,\epsilon}^{\delta_k}$ converge locally uniformly to a limiting conformal map $\varphi_{j,\epsilon}$. Since $(\mathcal{S}, x, y)$ almost surely does not have atoms, it follows that each of the limiting $\varphi_{j,\epsilon}$ with $j \leq |\mathcal{U}_\epsilon|$ satisfy $\varphi'_{j,\epsilon}(0) \neq 0$. This implies the result since, as mentioned above, the law of each of the $\mathcal{U}_\epsilon^{\delta_k}$ is the same as the law of each of the $\mathcal{U}_\epsilon$. □

Remark 6.4. Suppose that Y_t is a 3/2-stable Lévy process with only positive jumps and that Λ is a Poisson point process on $[0, t] \times \mathbf{R}_+$ with intensity measure $cds \otimes u^{-5/2}du$ where ds, du both denote Lebesgue measure and $c > 0$ is a constant. Then there exists a value of $c > 0$ such that Λ is equal in distribution to the set which consists of the pairs (t, u) where t is the time at which Y makes a jump and u is the size of the jump. This implies that if we observe only the jumps made by Y up to a random time t, then we can determine t by counting the number of jumps that Y has made with size between e^{-j-1} and e^{-j}, dividing by the factor $c_0 e^{3j/2}$ where $c_0 = \frac{2c}{3}(e^{3/2} - 1)$, and then sending $j \to \infty$. Using the same principle, we can almost surely determine the length of time that the time-reversal of a 3/2-stable Lévy excursion $e \colon [0, T] \to \mathbf{R}_+$ has been run if we only observe its jumps. Combining this with Proposition 6.3, this allows us to make sense of the quantum natural time parameterization of Γ where the jumps are provided by the quantum boundary lengths of the bubbles cut off by Γ.

Building on Remark 6.4, we have that the ordered collection of components cut off by a QLE$(8/3, 0)$ from its target point almost surely determines a 3/2-stable Lévy excursion. We will now show that the time-reversal of this excursion is almost surely equal to the process which gives the boundary length of Γ when parameterized using the quantum natural time parameterization (in the same way that the ordered sequence of bubbles cut off by an SLE_6 almost surely determines the evolution of the quantum length of

its outer boundary) and that the conditional law of the region (viewed as a quantum surface) given its quantum boundary length which contains the target point is the same as for an SLE_6 — a quantum disk with the given boundary length weighted by its quantum area.

Proposition 6.5. *Suppose that $(\mathcal{S}, x, y)$ is a doubly-marked quantum sphere with distribution $\mathsf{M}^2_{\mathrm{SPH}}$ and that Γ is a $\mathrm{QLE}(8/3, 0)$ on $\mathcal{S}$ from x to y. We take Γ to be parameterized by quantum natural time as described in Remark 6.4. For each fixed t, the conditional law of the (unique) complementary component of Γ_t viewed as a quantum surface given its boundary length is equal to the conditional law of the complementary component containing y of a whole-plane SLE_6 on $\mathcal{S}$ from x to y viewed as a quantum surface given its boundary length. That is, it is given by a quantum disk with the given boundary length weighted by its area. As t varies, the quantum boundary length of the complementary component of Γ_t evolves in the same manner as for a whole-plane SLE_6 (i.e., the time-reversal of a 3/2-stable Lévy excursion with only upward jumps) and is equal to the time-reversal of the Lévy excursion whose ordered sequence of jumps is given by the boundary lengths of the components swallowed by Γ.*

Proof. We shall assume that $(\mathcal{S}, x, y) = (\mathbf{C}, h, 0, \infty)$ and that Γ is parameterized by capacity as in the construction of the subsequential limits in the beginning of this subsection. Let (δ_k) be a sequence as in the beginning of this section and, for each k, let Γ^{δ_k} be the δ_k-approximation to $\mathrm{QLE}(8/3, 0)$. We assume that each of the Γ^{δ_k} are parameterized by capacity. For each $\epsilon > 0$, we let $\tau^{\delta_k}_{\epsilon,j}$ (resp. $\tau_{\epsilon,j}$) be the first capacity time t that Γ^{δ_k} (resp. Γ) has swallowed exactly j bubbles with quantum boundary length at least ϵ. Then we know that the joint law of the quantum surfaces parameterized by $\mathbf{C} \setminus \Gamma^{\delta_k}_{\tau^{\delta_k}_{\epsilon,j}}$ as j varies is equal in distribution to the corresponding family of surfaces in the case of an SLE_6 exploration of a doubly-marked quantum sphere sampled from $\mathsf{M}^2_{\mathrm{SPH}}$. Repeating the argument of Proposition 6.3 (i.e., passing to a further subsequence and recoupling the laws of the δ_k-approximations using the Skorohod embedding theorem), we can construct a coupling with an SLE_6 exploration of a doubly-marked quantum sphere so that the unexplored region for the $\mathrm{QLE}(8/3, 0)$ at these times is equal to the unexplored region at the corresponding time for an SLE_6. By passing to a further (diagonal) subsequence if necessary, we can construct a coupling so that this holds for all $\epsilon > 0$ simultaneously. That is, the joint law of the unexplored region for a $\mathrm{QLE}(8/3, 0)$ stopping times of the form $\tau_{\epsilon,j}$ is the same as for an SLE_6 exploration. Moreover, we can arrange so that the ordered sequence of bubbles cut off by Γ are equal to the sequence of bubbles cut off by the SLE_6. This implies the result because the stopping times of the form $\tau_{\epsilon,j}$ are dense in the quantum natural time parameterization. □

Proposition 6.6. *Suppose that we have the same setup as in Proposition 6.5 where we have taken $(\mathcal{S}, x, y) = (\mathbf{C}, h, 0, \infty)$. Then for each time t, we have that $\mathbf{C} \setminus \Gamma_t$ is a Hölder domain for each $t > 0$ fixed. In particular, $\partial\Gamma_t$ is almost surely conformally removable.*

Proof. This follows because the law of the surface parameterized by $\mathbf{C} \setminus \Gamma_t$ has the same law as the corresponding surface for an SLE_6, hence we can apply the argument of Proposition 5.12. □

6.3 Time parameterizations

In this article, we consider three different time parameterizations for $\mathrm{QLE}(8/3, 0)$:

1. Capacity time,
2. Quantum natural time, and
3. Quantum distance.

We have already introduced the first two parameterizations. We will now give the definition of the third. Suppose that Γ is a $\mathrm{QLE}(8/3, 0)$ parameterized by quantum natural time and let X_t be the quantum boundary length of the complementary component of Γ_t. We then set

$$s(t) = \int_0^t \frac{1}{X_u} du.$$

We refer to the time-parameterization given by $s(t)$ as the **quantum distance** parameterization for $\mathrm{QLE}(8/3, 0)$. The reason for this terminology is that, as we shall see in the next section, those points which are swallowed by Γ at quantum distance time d have distance d from x in the metric space that we construct. Let $\zeta = \inf\{t > 0 : X_t = 0\}$ and let

$$d_{\mathcal{Q}}(x, y) = \int_0^{\zeta} \frac{1}{X_u} du = s(\zeta). \tag{6.1}$$

Then $d_{\mathcal{Q}}(x, y)$ measures the quantum distance "from the left" of x and y (in the sense that x is the "left" argument of $d_{\mathcal{Q}}(x, y)$ and y is the "right" argument). One of the main inputs into the proof of Theorem 1.1 given in Section 8.2 is that $d_{\mathcal{Q}}(x, y)$ is symmetric in x and y and in fact almost surely determined by the underlying surface $\mathcal{S}$.

Lemma 6.7. *Suppose that Γ is a $\mathrm{QLE}(8/3, 0)$ with either the quantum distance or quantum natural time parameterization. We almost surely have for all $0 \leq s < t$ that Γ_s is a strict subset of Γ_t.*

Proof. This follows because it is almost surely the case that in each open interval of time in the quantum natural time parameterization Γ swallows a bubble and the same is also true for the quantum distance parameterization. □

Remark 6.8. Lemma 6.7 implies that the function which converts from the quantum natural time (resp. distance) parameterization to the capacity time parameterization is almost surely strictly increasing. It does not, however, rule out the possibility that this time change has jumps.

7 Symmetry

The purpose of this section is to establish an analog of Theorem 5.1 for QLE$(8/3, 0)$, stated as Theorem 7.1 below. This will be a critical ingredient for our proof of Theorem 1.1 given in Section 8. We begin by introducing a QLE$(8/3, 0)$ analog of the measure $\mathsf{M}^2_{\mathrm{SPH,D}}$ from Section 4.2. To define this measure, we let (δ_k) and $(\overline{\delta}_k)$ be two sequences as in Section 6.2 along which the δ-approximations to QLE$(8/3, 0)$ converge. We do not assume that (δ_k) and $(\overline{\delta}_k)$ are the same. We then let Θ be the measure on quadruples consisting of a doubly-marked quantum sphere $(\mathcal{S}, x, y)$, QLE$(8/3, 0)$ growth processes Γ and $\overline{\Gamma}$ from x to y (resp. y to x), and $U \in [0, 1]$. We assume that Γ (resp. $\overline{\Gamma}$) is produced from the limiting law associated with (δ_k) (resp. $(\overline{\delta}_k)$). A sample from Θ is produced by:

1. Picking $(\mathcal{S}, x, y)$ from $\mathsf{M}^2_{\mathrm{SPH}}$,
2. Picking Γ and $\overline{\Gamma}$ conditionally independently given $(\mathcal{S}, x, y)$, and
3. Taking U to be uniform in $[0, 1]$ independently of everything else.

We assume that both Γ and $\overline{\Gamma}$ have the quantum distance parameterization. Let $d_{\mathcal{Q}}(x, y)$ be as in (6.1) for Γ and $\overline{d}_{\mathcal{Q}}(y, x)$ be as in (6.1) for $\overline{\Gamma}$. We note that it is not *a priori* clear that the quantities $d_{\mathcal{Q}}(x, y)$ and $\overline{d}_{\mathcal{Q}}(y, x)$ should be related because the former is a measurement of distance between x and y from the "left" while the latter measures distance from the "right." Moreover, $d_{\mathcal{Q}}(x, y)$ and $\overline{d}_{\mathcal{Q}}(y, x)$ are defined with what *a priori* might be different subsequential limits of QLE$(8/3, 0)$.

We also let $\tau = U d_{\mathcal{Q}}(x, y)$, $\overline{\sigma} = U \overline{d}_{\mathcal{Q}}(y, x)$,

$$\overline{\tau} = \inf\{t \geq 0 : \overline{\Gamma}_t \cap \Gamma_\tau \neq \emptyset\}, \quad \text{and} \quad \sigma = \inf\{t \geq 0 : \Gamma_t \cap \overline{\Gamma}_{\overline{\sigma}} \neq \emptyset\}.$$

Note that τ (resp. $\overline{\sigma}$) is uniform in $[0, d_{\mathcal{Q}}(x, y)]$ (resp. $[0, \overline{d}_{\mathcal{Q}}(y, x)]$). We define measures $\Theta^{x \to y}$ and $\Theta^{y \to x}$ by setting

$$\frac{d\Theta^{x \to y}}{d\Theta} = d_{\mathcal{Q}}(x, y) \quad \text{and} \quad \frac{d\Theta^{y \to x}}{d\Theta} = \overline{d}_{\mathcal{Q}}(y, x). \tag{7.1}$$

Theorem 7.1. *Suppose that we have the setup described just above. Then the $\Theta^{x \to y}$ distribution of $\Gamma|_{[0,\tau]}$, $\overline{\Gamma}|_{[0,\overline{\tau}]}$,$(\mathcal{S}, x, y)$ is equal to the $\Theta^{y \to x}$ distribution of $\Gamma|_{[0,\sigma]}$, $\overline{\Gamma}|_{[0,\overline{\sigma}]}$, $(\mathcal{S}, x, y)$.*

7.1 Conditional independence in the limit

We will need to have a version of the following statement in the proof of Theorem 7.1. Suppose that we start with a doubly-marked quantum sphere $(\mathcal{S}, x, y)$ together with a



Figure 7.1: Illustration of the argument used to prove Theorem 7.1. The top shows the configuration of paths as in Figure 5.4. By Theorem 5.1, we know that the distribution of the path-decorated quantum surfaces parameterized by the light blue and red regions on the top and marked by the tip of the corresponding SLE_6 segments is symmetric under $\mathsf{M}^2_{\mathrm{SPH,D}}$. Reshuffling the two SLE_6 processes on the top yields the pair of $\mathrm{QLE}(8/3, 0)$ growths on the bottom. We will deduce our symmetry result for the surfaces on the bottom from the symmetry of the surfaces on the top. Part of this involves arguing that the asymmetry in the top which arises from knowing that $\overline{\eta}'(\bar{\mathfrak{t}}) \in \eta'([0, \mathfrak{t}])$ disappears in the limiting procedure used to construct $\mathrm{QLE}(8/3, 0)$.

δ_k-approximation of $\mathrm{QLE}(8/3, 0)$ from x to y and a $\overline{\delta}_k$-approximation of $\mathrm{QLE}(8/3, 0)$ from y to x, taken to be conditionally independent given $(\mathcal{S}, x, y)$. Then in the weak limit along sequences (δ_k) and $(\overline{\delta}_k)$ as in Section 6.2 we have that the resulting $\mathrm{QLE}(8/3, 0)$'s are conditionally independent given the underlying quantum surface, at least when the processes are grown up to a time at or before they first touch.

Proposition 7.2. *Suppose that (δ_k), $(\overline{\delta}_k)$ are two sequences as in Section 6.2. Suppose further that, for each k, we have a triple consisting of a doubly-marked quantum sphere $(\mathcal{S}^k, x^k, y^k) = (\mathbf{C}, h^k, 0, \infty)$, a δ_k-approximation Γ^k of $\mathrm{QLE}(8/3, 0)$ from 0 to ∞, and a $\overline{\delta}_k$-approximation $\overline{\Gamma}^k$ of $\mathrm{QLE}(8/3, 0)$ from ∞ to 0. We assume further that Γ^k and*

$\overline{\Gamma}^k$ *are conditionally independent given* h^k *and that both are parameterized by capacity as seen from their target point. Let* $(\mathcal{S}, x, y) = (\mathbf{C}, h, 0, \infty)$, Γ, $\overline{\Gamma}$ *be distributed as the weak limit of the aforementioned triple (so both of the limiting* $\mathrm{QLE}(8/3, 0)$*'s are parameterized by capacity as seen from their target point). Let* τ *be any stopping time for the filtration generated by* Γ *and the restriction of* h *to the interior of* Γ_t *so that* Γ_τ *is almost surely bounded and let* $\overline{\tau}$ *be the first time that* $\overline{\Gamma}$ *hits* Γ_τ*. Then the joint distribution of* $(\mathbf{C}, h, 0, \infty)$, $\Gamma|_{[0,\tau]}$, *and* $\overline{\Gamma}|_{[0,\overline{\tau}]}$ *is equal to the corresponding distribution if we had taken* Γ *and* $\overline{\Gamma}$ *to be* $\mathrm{QLE}(8/3, 0)$ *processes generated respectively from the limiting law associated with* (δ_k) *and* $(\overline{\delta}_k)$ *sampled conditionally independently given* h*.*

In order to establish Proposition 7.2, we will need to control the Radon-Nikodym derivative of the conditional law of the approximation of one of the $\mathrm{QLE}(8/3, 0)$'s given the other. It is not hard to find a counterexample which shows that if a sequence (X_k, Y_k, Z_k) converges weakly to (X, Y, Z) such that X_k, Y_k are conditionally independent given Z_k for each k then it is not necessarily true that X, Y are conditionally independent given Z. We begin with Lemma 7.3 which gives a condition under which the conditional independence of X_k, Y_k given Z_k implies the conditional independence of X, Y given Z. We will then use this condition in Lemma 7.5 to get the conditional independence of the limiting $\mathrm{QLE}(8/3, 0)$'s stopped upon exiting a pair of disjoint open sets from which Proposition 7.2 will follow.

Lemma 7.3. *Suppose that* (X_k, Y_k, Z_k) *is a sequence of random variables such that* X_k *and* Y_k *are conditionally independent given* Z_k *for each* k *and* $(X_k, Y_k, Z_k) \to (X, Y, Z)$ *weakly as* $k \to \infty$*. For each* k*, let* f_k *(resp.* g_k*) be the Radon-Nikodym derivative of the conditional law of* X_k *(resp.* Y_k*) given* Z_k *with respect to its unconditioned law. For each* k*, we let* m_k^X, m_k^Y, m_k^Z *respectively denote the laws of* X_k, Y_k, Z_k*. Assume that* $f_k = f$ *and* $g_k = g$ *do not depend on* k*, that* f, g *are continuous, and that* $f(X_k, Z_k)g(Y_k, Z_k)$ *is uniformly integrable for* $dm_k^X dm_k^Y dm_k^Z$*. Then* X, Y *are conditionally independent given* Z*.*

Proof. For simplicity we will prove the result in the case that f, g are bounded. The result in the general case follows from a simple truncation argument. Let m^X, m^Y, m^Z be the respective laws of X, Y, Z. Then the joint law of (X_k, Y_k, Z_k) is given by

$$f(X_k, Z_k)g(Y_k, Z_k)dm_k^X dm_k^Y dm_k^Z.$$

Suppose that F is a bounded, continuous function of (X_k, Y_k, Z_k). Then the weak convergence of (X_k, Y_k, Z_k) to (X, Y, Z) as $k \to \infty$ implies that

$$\begin{aligned} &\int F(X_k, Y_k, Z_k)f(X_k, Z_k)g(Y_k, Z_k)dm_k^X dm_k^Y dm_k^Z \\ \to &\int F(X, Y, Z)f(X, Z)g(Y, Z)dm^X dm^Y dm^Z \quad \text{as} \quad k \to \infty. \end{aligned}$$

Therefore $f(X,Z)g(Y,Z)dm^X dm^Y dm^Z$ gives the joint law of (X,Y,Z). From the form of the joint law, it is clear that X, Y are conditionally independent given Z, as desired. □

In order to be able to apply Lemma 7.3, we need to give the Radon-Nikodym derivative of the conditional law of a quantum sphere in a region conditional on its realization in a disjoint region. We will state this result in the setting of the infinite measure $\mathsf{M}_{\mathrm{BES}}$ on doubly-marked quantum spheres described in [DMS14, MS15b] constructed using the excursion measure associated with a Bessel process because this will allow us to give the simplest formulation of the result. We recall [MS15b, Theorem 1.4] which states that there exists a constant $c_{\mathrm{LB}} > 0$ such that $\mathsf{M}^2_{\mathrm{SPH}} = c_{\mathrm{LB}}\mathsf{M}_{\mathrm{BES}}$. It will be convenient in what follows to produce samples of quantum spheres using $\mathsf{M}_{\mathrm{BES}}$ rather than $\mathsf{M}^2_{\mathrm{SPH}}$ because the Bessel description is more amenable to using the Markov property for the GFF. We will briefly recall this construction; see the introduction of [MS15b] as well as [DMS14, Section 4.4 and Section 5.3] for additional detail. We let $\mathcal{Q} = \mathbf{R} \times [0, 2\pi]$ with the top and bottom identified be the infinite cylinder. We then let $\mathcal{H}(\mathcal{Q})$ be the Hilbert space closure of $C_0^\infty(\mathcal{Q})$ with respect to the Dirichlet inner product

$$(f,g)_\nabla = \frac{1}{2\pi}\int \nabla f(x)\cdot\nabla g(x)dx \tag{7.2}$$

and we let $\mathcal{H}_1(\mathcal{Q})$ (resp. $\mathcal{H}_2(\mathcal{Q})$) be the subspace of $\mathcal{H}(\mathcal{Q})$ consisting of those functions which are constant (resp. have mean zero) on vertical lines. Then $\mathcal{H}_1(\mathcal{Q}) \oplus \mathcal{H}_2(\mathcal{Q})$ gives an orthogonal decomposition of $\mathcal{H}(\mathcal{Q})$ [DMS14, Lemma 4.2]. A sample can be produced from $\mathsf{M}_{\mathrm{BES}}$ as follows.

- Take the projection of h onto $\mathcal{H}_1(\mathcal{Q})$ to be given by $\frac{2}{\gamma}\log Z$ reparameterized to have quadratic variation du where Z is picked from the Itô excursion measure $\nu_\delta^{\mathrm{BES}}$ of a Bessel process of dimension $\delta = 4 - \frac{8}{\gamma^2}$. This defines Z modulo horizontal translation.
- Sample the projection onto $\mathcal{H}_2(\mathcal{Q})$ independently from the law of the corresponding projection of a whole-plane GFF on $\mathcal{Q}$.

(See [MS15b, Section 2.1.1] for a reminder of the construction of $\nu_\delta^{\mathrm{BES}}$.) Throughout, we let $\mathcal{Q}_\pm \subseteq \mathcal{Q}$ be the half-infinite cylinders given by $[0, 2\pi] \times \mathbf{R}_\pm$ with the top and bottom identified.

Lemma 7.4. *Fix $r \in \mathbf{R}$. Suppose that $(\mathcal{Q}, h, -\infty, +\infty)$ is sampled from $\mathsf{M}_{\mathrm{BES}}$ conditioned on the supremum of the projection of h onto $\mathcal{H}_1(\mathcal{Q})$ exceeding r (this defines a probability measure). We take the horizontal translation so that the projection of h onto $\mathcal{H}_1(\mathcal{Q})$ first hits r at $u = 0$. Let $U \subseteq \mathcal{Q}$ be a neighborhood of $+\infty$ which is contained in $\mathcal{Q}_+$ and let V be a neighborhood of $\mathcal{Q}_-$ such that $\mathrm{dist}(U, V) > 0$. Let $\mathcal{Z}_{U,V}$ be defined by*

$$\mathcal{Z}_{U,V} = Z_{U,V}^{-1}\exp\left((h|_U, g|_U)_\nabla\right) \tag{7.3}$$

where g is equal to the harmonic extension of $f - h|_V$ from V to $\mathcal{Q} \setminus V$ and $Z_{U,V}$ is a normalizing constant. Then the law of a sample produced from the law of h weighted by $\mathcal{Z}_{U,V}$ and then restricted to U is equal to the law of the restriction $h|_U$ of h to U conditional on the restriction $h|_V$ of h to V being equal to f.

Proof. We first recall that if h is a GFF on a domain $D \subseteq \mathbf{C}$ and $f \in H(D)$ then the Radon-Nikodym derivative of the law of $h + f$ with respect to the law of h is given by $Z^{-1} \exp((h, f)_\nabla)$ where Z is a normalization constant. (This is proved by using that the Radon-Nikodym derivative of the law of a $N(\mu, 1)$ random variable with respect to the law of a $N(0, 1)$ random variable is given by $Z^{-1} e^{x\mu}$ where Z is a normalizing constant.)

In order to make use of the aforementioned fact in the setting of the lemma, we first note that from the construction of $\mathsf{M}_{\mathrm{BES}}$ it follows that the conditional law of $h|_U$ given $h|_V$ is equal to that of a GFF on $\mathcal{Q} \setminus V$ restricted to U with Dirichlet boundary conditions on ∂V and at $+\infty$ given by those of h. Consequently, by the Markov property for the GFF, we can sample from the law of $h|_U$ conditioned on $h|_V = f$ by:

1. Sampling h according to its unconditioned law and then
2. Adding to $h|_U$ the harmonic extension g of $f - h|_V$ from V to $\mathcal{Q} \setminus V$.

We can extract from this the result as follows. We let $\widetilde{g} = g\phi$ where $\phi \in C^\infty(\mathcal{Q})$ is such that $\phi \equiv 0$ in a neighborhood of V and $\phi|_U \equiv 1$. We note that $\widetilde{g}$ is measurable with respect to $\sigma(h|_V)$. Then we have that $(h + \widetilde{g})|_U$ has the law of $h|_U$ given $h|_V \equiv f$. Moreover, with $W = \mathcal{Q} \setminus (U \cup V)$ we have that the Radon-Nikodym derivative of the conditional law of $h + \widetilde{g}$ given $h|_V$ with respect to the conditional law of h given $h|_V$ is equal to

$$\widetilde{Z}^{-1} \exp((h, \widetilde{g})_\nabla) = \widetilde{Z}^{-1} \exp((h|_W, \widetilde{g}|_W)_\nabla + (h|_U, g|_U)_\nabla) \tag{7.4}$$

where $\widetilde{Z}$ is a normalizing constant and we used that $\widetilde{g}|_U \equiv g|_U$. Integrating over the randomness of the first summand in the exponential on the right side of (7.4) and using the conditional independence of $(h|_W, \widetilde{g}|_W)_\nabla$ and $(h|_U, g|_U)_\nabla$ given $h|_V$ implies the result. (Recall that for test functions $\psi, \widetilde{\psi}$ we have that the covariance of $(h, \psi)_\nabla$ and $(h, \widetilde{\psi})_\nabla$ is given by $(\psi, \widetilde{\psi})_\nabla$ so that the two random variables are independent in the case that the supports of ψ, $\widetilde{\psi}$ are disjoint.) □

Lemma 7.5. *Suppose that $(\mathcal{S}, x, y) = (\mathbf{C}, h, 0, \infty)$ is as in Lemma 7.4 where the embedding is given by applying the change of coordinates $\mathcal{Q} \to \mathbf{C}$ given by $z \mapsto e^z$. Let $V \subseteq \mathbf{C}$ be a bounded open neighborhood of $\mathbf{D}$ and $U \subseteq \mathbf{C}$ be an open neighborhood of ∞ such that $\mathrm{dist}(U, V) > 0$. Let (δ_k), $(\overline{\delta}_k)$ be sequences as in Section 6.2. Suppose that Γ^k is a δ_k-approximation to $\mathrm{QLE}(8/3, 0)$ from 0 to ∞ and $\overline{\Gamma}^k$ is a $\overline{\delta}_k$-approximation to $\mathrm{QLE}(8/3, 0)$ from ∞ to 0. Assume that Γ^k and $\overline{\Gamma}^k$ are conditionally independent given h and that both have the capacity time parameterization as seen from their target*

point. Let τ_V (resp. τ_U) be the infimum of times t that Γ^k (resp. $\overline{\Gamma}^k$) is not in V (resp. U). Then the joint law of $(\mathbf{C}, h, 0, \infty)$, $\Gamma^k|_{[0,\tau_V]}$, $\overline{\Gamma}^k|_{[0,\tau_U]}$ converges weakly as $k \to \infty$ to a triple which consists of a doubly-marked quantum sphere $(\mathbf{C}, h, 0, \infty)$ conditioned as in Lemma 7.4, a $\mathrm{QLE}(8/3, 0)$ process Γ from 0 to ∞ stopped upon exiting V, and a $\mathrm{QLE}(8/3, 0)$ process $\overline{\Gamma}$ from ∞ to 0 stopped upon exiting U with both Γ and $\overline{\Gamma}$ parameterized by capacity as seen from their target point. Moreover, Γ and $\overline{\Gamma}$ are conditionally independent given h.

Proof. We know that the joint law of the restriction $h|_V$ of h to V along with Γ^k converges weakly as $k \to \infty$. We likewise know that the same is true for $h|_U$ along with $\overline{\Gamma}^k$. As mentioned earlier, by [MS15b, Theorem 1.4], the Radon-Nikodym derivative of $\mathsf{M}^2_{\mathrm{SPH}}$ with respect to $\mathsf{M}_{\mathrm{BES}}$ is given by a constant. It therefore follows that we also have weak convergence as $k \to \infty$ when we produce our sample of $(\mathcal{S}, x, y)$ using $\mathsf{M}_{\mathrm{BES}}$ in place of $\mathsf{M}^2_{\mathrm{SPH}}$ and it suffices to show that the weak limit which comes from $\mathsf{M}_{\mathrm{BES}}$ has the desired conditional independence property. So, in what follows, we assume that $(\mathcal{S}, x, y)$ is sampled from $\mathsf{M}_{\mathrm{BES}}$.

By the locality property for SLE_6, observe that the Radon-Nikodym derivative $\mathcal{Z}_{U,V}$ of the conditional law of $h|_U$ and $\overline{\Gamma}^k|_{[0,\tau_U]}$ given $h|_V$ and $\Gamma^k|_{[0,\tau_V]}$ with respect to the unconditioned law is equal to the Radon-Nikodym derivative of the conditional law of $h|_U$ given $h|_V$ with respect to the unconditioned law of $h|_U$. By the explicit form of $\mathcal{Z}_{U,V}$ given in (7.3) of Lemma 7.4, we see that $\mathcal{Z}_{U,V}$ is a continuous function of the harmonic extension of $h|_V$ to $\mathbf{C} \setminus V$ restricted to U. By passing to subsequences of (δ_k) and $(\overline{\delta}_k)$ if necessary, we may assume that this harmonic extension converges weakly with along the other previously mentioned variables $k \to \infty$. Combining everything with Lemma 7.3 implies the result. □

Proof of Proposition 7.2. Lemma 7.5 implies that the assertion of the proposition holds if we replace τ with the first time t that Γ exits a bounded open neighborhood V of 0 and we replace $\overline{\tau}$ with the first time t that $\overline{\Gamma}$ exits a neighborhood U of ∞ with $\mathrm{dist}(U, V) > 0$. By taking a limit as U increases to the interior of the complement of V, it follows that the assertion of the lemma holds if we take τ to be the first time t that Γ exits V and $\overline{\tau}$ to be the first time t that $\overline{\Gamma}$ exits the interior of $\mathbf{C} \setminus V$. The result follows because this holds for all V which are bounded and open. □

7.2 Reshuffling a pair of SLE_6's

7.2.1 Setup

We begin by describing the setup and notation that we will use throughout this subsection. We first suppose that $\mathcal{X}$, $\overline{\mathcal{X}}$, $\mathcal{B}$, η', $\overline{\eta}'$, $\mathfrak{t}$, and $\bar{\mathfrak{t}}$ are as in Theorem 5.1. (In the setting of the statement of Theorem 7.1, $\mathfrak{t}$ plays the role of τ and $\bar{\mathfrak{t}}$ plays the

role of $\overline{\tau}$.) Then we know that the $\mathsf{M}^2_{\mathrm{SPH,D}}$ distribution of $(\mathcal{X}, \overline{\mathcal{X}}, \mathcal{B})$ is invariant under the operation of swapping $\mathcal{X}$ and $\overline{\mathcal{X}}$. Let (δ_k), $(\overline{\delta}_k)$ be sequences of positive numbers decreasing to 0 as in Section 6.2. Let $\Gamma^k|_{[0,\mathfrak{t}]}$ be the δ_k-approximation to QLE$(8/3, 0)$ which we take to be coupled with $\eta'|_{[0,\mathfrak{t}]}$ so that the bubbles that it separates from its target are the same as quantum surfaces as the bubbles separated by $\eta'|_{[0,\mathfrak{t}]}$ from its target point. That is, in the construction of $\Gamma^k|_{[0,\mathfrak{t}]}$ we take the time-reversal of the Lévy excursion e to be the same as the time-reversal of the Lévy excursion for η' up to time $\mathfrak{t}$ and we take the bubbles that $\Gamma^k|_{[0,\mathfrak{t}]}$ separates from its target point up to time $\mathfrak{t}$ to be the same as the bubbles that η' separates from its target point up to time $\mathfrak{t}$. We sample the rest of the process and quantum sphere conditionally independently given its realization of up to time $\mathfrak{t}$.

We likewise let $\overline{\Gamma}^k|_{[0,\overline{\mathfrak{t}}]}$ be the $\overline{\delta}_k$-approximation to QLE$(8/3, 0)$ which we take to be coupled with $\overline{\eta}'|_{[0,\overline{\mathfrak{t}}]}$ so that the bubbles it separates from its target are the same as quantum surfaces as the bubbles separated by $\overline{\eta}'|_{[0,\overline{\mathfrak{t}}]}$ from its target point. We sample the rest of the process and the quantum sphere on which it is growing conditionally independently given its realization up to time $\overline{\mathfrak{t}}$.

Consider the doubly-marked quantum sphere $(\mathcal{S}^k, x^k, y^k)$ which arises by starting with the doubly-marked sphere $(\mathcal{S}, x, y)$ as above, cutting out the quantum surfaces separated by $\eta'|_{[0,\mathfrak{t}]}$ and $\overline{\eta}'|_{[0,\overline{\mathfrak{t}}]}$ from their respective target points, and then gluing in according to quantum boundary length the corresponding quantum surfaces for $\Gamma^k|_{[0,\mathfrak{t}]}$ and $\overline{\Gamma}^k|_{[0,\overline{\mathfrak{t}}]}$. (See Figure 7.1 for an illustration.) We identify the tip of the final SLE_6 segment used to build $\Gamma^k|_{[0,\mathfrak{t}]}$ with $\eta'(\mathfrak{t})$ and likewise for $\overline{\Gamma}^k|_{[0,\overline{\mathfrak{t}}]}$ and $\overline{\eta}'(\overline{\mathfrak{t}})$. Proposition 5.12 implies that the resulting quantum surface is uniquely determined by this gluing operation. Moreover, the distribution of $(\mathcal{S}^k, x^k, y^k)$ is equal to that of $(\mathcal{S}, x, y)$. We abuse notation and let Γ^k and $\overline{\Gamma}^k$ be the resulting δ_k and $\overline{\delta}_k$-approximations to QLE$(8/3, 0)$ and let $\mathcal{B}^k$ be the region in $\mathcal{S}^k$ which is not in either $\Gamma^k_{\mathfrak{t}}$ or $\overline{\Gamma}^k_{\overline{\mathfrak{t}}}$.

Throughout, we will write τ (resp. $\overline{\tau}$) for $\mathfrak{t}$ (resp. $\overline{\mathfrak{t}}$) when referring to either Γ^k or Γ (resp. $\overline{\Gamma}^k$ or $\overline{\Gamma}$) and use $\mathfrak{t}$ (resp. $\overline{\mathfrak{t}}$) when referring to η' (resp. $\overline{\eta}'$).

We let Θ_k denote the joint distribution of $(\mathcal{S}^k, x^k, y^k)$, Γ^k, $\overline{\Gamma}^k$, τ and $\overline{\tau}$ and let $\Theta_k^{x\to y}$ be given by Θ_k weighted by the amount of time it takes Γ to reach its target point. We define $\Theta_k^{y\to x}$ similarly except we weight by the amount of time it takes $\overline{\Gamma}$ to reach its target point and we write $\overline{\sigma}$ for uniform variable between 0 and this time and let σ be the first time that Γ^k_t hits $\overline{\Gamma}^k_{\overline{\sigma}}$. We write $\overline{\mathcal{B}}^k$ for the surface which is the complement of Γ^k_σ and $\overline{\Gamma}^k_{\overline{\sigma}}$. Note that $\Theta_k^{x\to y}$ and $\Theta_k^{y\to x}$ are the analogs of $\Theta^{x\to y}$ and $\Theta^{y\to x}$ defined at the beginning of this section except we have used the approximations to QLE$(8/3, 0)$ instead of the limits.

Let $(\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B})$ be the configuration on the doubly-marked quantum sphere $(\mathcal{S}, x, y)$ which arises by taking the limit of $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{B}^k)$ under $\Theta_k^{x\to y}$ as $k \to \infty$. This limit exists by Proposition 7.2 and the distribution of $(\mathcal{S}, x, y)$, $\Gamma|_{[0,\tau]}$, and $\overline{\Gamma}|_{[0,\overline{\tau}]}$ as

constructed is equal to its distribution under $\Theta^{x\to y}$. We likewise let $(\Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}})$ be the configuration on the doubly-marked quantum sphere $(\mathcal{S}, x, y)$ which arises by taking the limit of $(\Gamma^k|_{[0,\sigma]}, \overline{\Gamma}^k|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}}^k)$ under $\Theta_k^{y\to x}$ as $k \to \infty$. Analogously, we can construct the law of $(\mathcal{S}, x, y)$, $\Gamma|_{[0,\sigma]}$, $\overline{\Gamma}|_{[0,\overline{\sigma}]}$ under $\Theta^{y\to x}$ as the same type of limit.

7.2.2 Structure of the meeting QLEs

In this section we will prove two results regarding the structure of the triple $\Gamma|_{[0,\tau]}$, $\overline{\Gamma}|_{[0,\overline{\tau}]}$, and $\mathcal{B}$ under $\Theta^{x\to y}$ (as well as analogous results in the setting of $\Theta^{y\to x}$).

Lemma 7.6. *Under $\Theta^{x\to y}$, we have that $\Gamma|_{[0,\tau]}$ and $\overline{\Gamma}|_{[0,\overline{\tau}]}$ almost surely intersect at a single point. In particular, the region $\mathcal{B}$ of $\mathcal{S}$ outside of $\Gamma|_{[0,\tau]}$ and $\overline{\Gamma}|_{[0,\overline{\tau}]}$ is almost surely connected. Moreover, the conditional law of $\mathcal{B}$ given its boundary length is that of a quantum disk and $\partial\mathcal{B}$ is almost surely conformally removable in any embedding of $\mathcal{S}$ into $\mathbf{C}$. The same holds in the setting of $\Theta^{y\to x}$ in place of $\Theta^{x\to y}$.*

See Figure 7.2 for an illustration of the setup and proof of Lemma 7.6.

Proof of Lemma 7.6. We know that the statement of the lemma holds in the setting of each of the δ_k-approximations in the proof of Theorem 7.1. Indeed, this follows because a radial SLE_6 segment almost surely hits the boundary of any fixed domain for the first time at a single point. For each $k \in \mathbf{N}$, we pick z^k in $\mathcal{B}^k$ according to the quantum area measure and then parameterize $\mathcal{B}^k$ by $(\mathbf{D}, h^k)$ by using the change of coordinates given by the unique conformal map $\varphi^k \colon \mathbf{D} \to \mathcal{B}^k$ with $\varphi^k(0) = z^k$ and $(\varphi^k)'(0) > 0$. We note that the distribution of $(\mathbf{D}, h^k)$ does not depend on k and that, by Proposition 5.12 its law conditionally on its boundary length is that of a quantum disk.

Fix $\ell > 0$ and assume that we are working on the event that the quantum boundary length of $\partial\Gamma_\tau^k$ is in $[\ell, \ell+1]$. Since this event has positive and finite $\Theta_k^{x\to y}$ mass, we can condition $\Theta_k^{x\to y}$ on it and we get a probability measure. Suppose for contradiction that there is a chance $p > 0$ that $\partial\Gamma_\tau \cap \partial\overline{\Gamma}_{\overline{\tau}}$ consists of more than a single point. This would imply that the following is true. Let u^k be the unique element of $\partial\Gamma_\tau^k \cap \partial\overline{\Gamma}_{\overline{\tau}}^k$. Then there exists $r > 0$ such that with probability at least $p/2$ there is a point v^k on $\partial\Gamma_\tau^k$ such that the quantum length of either the clockwise or counterclockwise segment of $\partial\Gamma_\tau^k$ from u^k to v^k is at least r and its harmonic measure as seen from z^k is $o(1)$ as $k \to \infty$. In particular, the arc length of the preimage of this interval under φ^k is $o(1)$ as $k \to \infty$. This implies that $(\mathbf{D}, h^k)$ converges as $k \to \infty$ to a field whose associated quantum boundary measure has a positive chance of having an atom on $\partial\mathbf{D}$. This is a contradiction because the law of $(\mathbf{D}, h^k)$ does not depend on k and its associated quantum boundary measure almost surely does not have an atom on $\partial\mathbf{D}$ (Proposition 5.12 and [DS11]). This proves the first assertion of the lemma.

The second assertion follows because the distribution of $\mathcal{B}^k$ does not depend on k and the final assertion follows by using the argument of Proposition 5.12 (i.e., any



Figure 7.2: Illustration of the setup and proof of Lemma 7.6. Shown on the left are two conditionally independent δ_k-approximations $\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}$ of QLE$(8/3, 0)$ on a doubly-marked quantum sphere $(\mathcal{S}^k, x^k, y^k)$ where $\Gamma^k|_{[0,\tau]}$ is grown up to a typical quantum distance time and $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ is grown up until the first time that it hits $\Gamma^k|_{[0,\tau]}$. It is almost surely the case that these two processes intersect at exactly one point because radial SLE_6 almost surely hits the boundary of any given simply connected domain for the first time at a single point; call the complementary region $\mathcal{B}^k$ and let z^k be a uniformly chosen point from the quantum measure on $\mathcal{B}^k$. By Proposition 5.12, $\mathcal{B}^k$ is a quantum disk conditionally on its boundary length and the distribution of $\mathcal{B}^k$ does not depend on k. If, in the limit as $k \to \infty$, there was a positive chance that the processes intersect at more than one point then there would be a positive chance that the δ_k-approximations would form a narrow fjord the part of whose boundary which is in $\partial\Gamma^k_\tau$ has uniformly positive quantum length. This leads to a contradiction because harmonic measure considerations imply that the inverse image of the green part of the fjord under the unique conformal map $\varphi^k \colon \mathbf{D} \to \mathcal{B}^k$ with $\varphi^k(0) = z^k$ and $(\varphi^k)'(0) > 0$ would be a segment with a uniformly positive amount of quantum length (indicated in green on the right) but with arc length tending to 0. In particular, it would converge in the $k \to \infty$ limit to an atom of the boundary measure.

embedding of a quantum disk into a quantum sphere almost surely has a conformally removable boundary). □

Lemma 7.7. *Under $\Theta^{x \to y}$, the point on $\partial\Gamma_\tau$ (resp. $\partial\overline{\Gamma}_{\overline{\tau}}$) which is glued to the almost surely unique pinch point of $\partial\mathcal{B}$ is uniformly distributed from the quantum boundary measure on $\partial\Gamma_\tau$ (resp. $\partial\overline{\Gamma}_{\overline{\tau}}$). The same holds in the setting of $\Theta^{y \to x}$ in place of $\Theta^{x \to y}$.*

Proof. It is easy to see that this is the case for $\Gamma|_{[0,\tau]}$ based on the construction. Let $[\xi, \overline{\tau}]$ be the interval of time in which $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ is drawing its final segment of SLE_6. The claim in the case of $\overline{\Gamma}|_{[0,\overline{\tau}]}$ follows because for each of the approximations $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ we know that the starting point of the final SLE_6 segment is uniformly distributed on $\partial\overline{\Gamma}^k_\xi$ and this final SLE_6 segment collapses to a point as $k \to \infty$. □

7.2.3 Preliminary symmetry statements

In order to work towards the proof of Theorem 7.1, we will now:

- Show that the $\Theta_k^{x\to y}$ distribution of $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{B}^k)$ is equal to the $\Theta_k^{y\to x}$ distribution of $(\Gamma^k|_{[0,\sigma]}, \overline{\Gamma}^k|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}}^k)$ (Lemma 7.8)
- Deduce from this that the $\Theta^{x\to y}$ distribution of $(\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B})$ is equal to the $\Theta^{y\to x}$ distribution of $(\Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}})$ (Lemma 7.10)

The proof of Theorem 7.1 will not require much additional work once we have established these lemmas.

Lemma 7.8. *The $\Theta_k^{x\to y}$ distribution of $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{B}^k)$ is the same as the $\Theta_k^{y\to x}$ distribution of $(\Gamma^k|_{[0,\sigma]}, \overline{\Gamma}^k|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}}^k)$.*

In the statement of Lemma 7.8, we view $\Gamma^k|_{[0,\tau]}$, $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$, $\Gamma^k|_{[0,\sigma]}$, $\overline{\Gamma}^k|_{[0,\overline{\sigma}]}$ as random variables taking values in the space of beaded quantum surfaces with a single marked point which corresponds to the tip of the final SLE_6 segment in the growth process and we view $\mathcal{B}^k, \overline{\mathcal{B}}^k$ as random variables taking values in the space of quantum surfaces.

Remark 7.9. As in Remark 5.2, we emphasize that Lemma 7.8 does not imply that the $\Theta_k^{x\to y}$ distribution of $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{S}^k)$ is equal to the $\Theta_k^{y\to x}$ distribution of $(\Gamma^k|_{[0,\sigma]}, \overline{\Gamma}^k|_{[0,\overline{\sigma}]}, \mathcal{S}^k)$ because of the asymmetry which arises since the tip of $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ is contained in $\partial\Gamma^k_\tau$ while the tip of $\Gamma^k|_{[0,\tau]}$ is uniformly distributed according to the quantum boundary measure on $\partial\Gamma^k_\tau$.

Proof of Lemma 7.8. The result is a consequence of the following three observations.

First, the ordered sequence of bubbles separated by $\Gamma^k|_{[0,\tau]}$ from its target point is equal to the ordered sequence of bubbles separated by $\eta'|_{[0,\mathfrak{t}]}$ from its target point and the same is true for $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ and $\overline{\eta}'|_{[0,\overline{\mathfrak{t}}]}$. Theorem 5.1 thus implies that the collection of δ_k-quantum natural time length SLE_6 segments (viewed as quantum surfaces) which make up $\Gamma^k|_{[0,\tau]}$ under $\Theta_k^{x\to y}$ have the same joint law as the corresponding collection for $\Gamma^k|_{[0,\sigma]}$ under $\Theta_k^{y\to x}$. It also implies that the quantum lengths of the outer boundary of

$\Gamma^k|_{[0,\tau]}$ at the reshuffling times $j\delta_k$ for $1 \leq j \leq \lfloor \tau/\delta_k \rfloor$ have the same joint law under $\Theta_k^{x\to y}$ as the quantum lengths of the outer boundary of $\Gamma^k|_{[0,\sigma]}$ at the reshuffling times $j\delta_k$ for $1 \leq j \leq \lfloor \sigma/\delta_k \rfloor$ under $\Theta_k^{y\to x}$. Indeed, this follows since these quantum boundary lengths are determined by the bubbles that the processes separate from their target points and the bubbles determine the SLE_6 segments.

Second, the starting locations of each of the SLE_6 segments which make up $\Gamma^k|_{[0,\tau]}$ have the same conditional law under $\Theta_k^{x\to y}$ given the bubbles as in the case of $\Gamma^k|_{[0,\sigma]}$ under $\Theta_k^{y\to x}$. Indeed, in both cases, these starting locations are distributed uniformly from the quantum boundary measure.

Both of the first two observations also apply for $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ under $\Theta_k^{x\to y}$ and $\overline{\Gamma}^k|_{[0,\overline{\sigma}]}$ under $\Theta_k^{y\to x}$.

Third, the previous two observations imply that the induced joint distribution of $\Gamma^k|_{[0,\tau]}$ and $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ under $\Theta_k^{x\to y}$ is the same as that of $\Gamma^k|_{[0,\sigma]}$ and $\overline{\Gamma}^k|_{[0,\overline{\sigma}]}$ under $\Theta_k^{y\to x}$. We also note that $\mathcal{B}^k$ is conditionally independent given its boundary length of $\Gamma^k|_{[0,\tau]}$ and $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$ under $\Theta_k^{x\to y}$. The same is also true under $\Theta^{y\to x}$, so combining implies the desired result. □

The next step is to show that the symmetry established in Lemma 7.8 also holds for $(\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B})$ under $\Theta^{x\to y}$ and $(\Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}})$ under $\Theta^{y\to x}$.

Before we state this result, we need to introduce a certain notion of equivalence for $\mathrm{QLE}(8/3, 0)$ processes. We recall from Proposition 6.5 that the complementary region of a $\mathrm{QLE}(8/3, 0)$ run up to a given quantum distance time has the same law as the corresponding region for an SLE_6 and from Proposition 6.6 that the boundary of a $\mathrm{QLE}(8/3, 0)$ can be parameterized by a continuous curve. This allows us to induce a quantum boundary length measure on the prime ends of the boundary of a $\mathrm{QLE}(8/3, 0)$ even though it is not *a priori* clear at this point whether this boundary measure can be defined in a way which is intrinsic to the $\mathrm{QLE}(8/3, 0)$. (At this point, we do know that the total boundary length is intrinsic to the $\mathrm{QLE}(8/3, 0)$ because it can be determined by boundary lengths of the bubbles it has cut off from its target point.) Throughout, we will say that two $\mathrm{QLE}(8/3, 0)$'s coupled on a common probability space and each stopped at a given time are equivalent if there exists a homeomorphism which takes one to the other which:

- Preserves the induced quantum boundary length on the prime ends of the boundary of the $\mathrm{QLE}(8/3, 0)$, and is
- Conformal in the interior of the $\mathrm{QLE}(8/3, 0)$.

As we will shall see later, this notion of equivalence is useful because it encodes the information necessary for the operation of welding according to quantum boundary length to be well-defined and uniquely determined (i.e., we will be in a setting in which we can apply the conformal removability of the boundary; recall Proposition 2.1).

Lemma 7.10. *The $\Theta^{x\to y}$ distribution of $(\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B})$ is equal to the $\Theta^{y\to x}$ distribution of $(\Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}})$ where we use the notion of equivalence for* QLE$(8/3, 0)$ *processes as described just above.*

We note that Lemma 7.10 is not a trivial consequence of Lemma 7.8. In order to prove the result, we will construct a coupling of the $\Theta^{x\to y}$ distribution of $(\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B})$ and the $\Theta^{y\to x}$ distribution of $(\Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}})$ on the same space so that there is a map of the above type which takes the former QLE$(8/3, 0)$'s to the latter. The non-trivial issue one has to deal with in order to push this argument through is that there is an asymmetry in the locations of the tips in the approximations to the former and latter QLE$(8/3, 0)$'s. Our approach to dealing with this asymmetry is to perform a "local surgery" near the tip and is illustrated in Figure 7.3 and Figure 7.4. We begin with the following observation.



Figure 7.3: Illustration of the setup for Lemma 7.11. Here, $[\xi, \tau]$ is the interval of time in which $\Gamma^k|_{[0,\tau]}$ is drawing its final segment of SLE_6 (shown in yellow). The purple region shows the final segment of SLE_6 for $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$. The green region shows the surface which is cut off by another $c + \xi - \tau$, $c > 0$ fixed and small, units of time by an SLE_6 starting from the tip of Γ at time τ so that the region $\mathcal{R}$ which is given by the union of the yellow and green regions corresponds to the region cut off by running an SLE_6 for c units of time.

Lemma 7.11. *Consider the law on triples of quantum surfaces $(\Upsilon^k, \overline{\Upsilon}^k, \widehat{\mathcal{B}}^k)$ which is defined as follows. We first sample $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{B}^k)$ from $\Theta^{x\to y}$. Let $\xi = \delta_k \lfloor \tau/\delta_k \rfloor$ be such that $[\xi, \tau]$ is the interval of time in which $\Gamma^k|_{[0,\tau]}$ is drawing its final segment of* SLE_6*, let $\eta'|_{[\xi,\infty)}$ be the corresponding* SLE_6 *process (parameterized by quantum distance time), and for $c > 0$ fixed and small, let $\zeta = \xi + c > \tau$. Let $\mathcal{R}$ be the beaded quantum surface which is cut out by $\eta'|_{[\xi,\zeta]}$. Fix $r \in \mathbf{R}$ and $\epsilon > 0$. On the event E that $\mathcal{R}$ is disjoint from $\overline{\Gamma}^k|_{[0,\overline{\tau}]}$, we resample $\mathcal{R}$ (with its overall boundary length fixed) and $\eta'|_{[\xi,\zeta]}$*

conditioned on the event $F_{r,\epsilon}$ that the difference in the quantum boundary length of $\partial\Gamma^k_\xi$ and $\partial\Gamma^k_\tau$ from r is at most ϵ. Let $(\widehat{\mathcal{S}}^k, \widehat{x}^k, \widehat{y}^k)$ be the resulting doubly-marked quantum sphere and $(\Upsilon^k, \overline{\Upsilon}^k, \widehat{\mathcal{B}}^k)$ be the resulting pair of $\mathrm{QLE}(8/3, 0)$ *growths together with their complementary region in $\widehat{\mathcal{B}}^k$. Then the joint law of $(\Upsilon^k, \overline{\Upsilon}^k, \widehat{\mathcal{B}}^k)$ and $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{B}^k)$ satisfies the following property. On E, there exists a map $\varphi^k \colon \mathcal{S}^k \to \widehat{\mathcal{S}}^k$ which is conformal outside of $\mathcal{R}$ and takes Γ^k_τ to Υ^k, $\overline{\Gamma}^k_{\overline{\tau}}$ to $\overline{\Upsilon}^k$, and $\mathcal{B}^k$ to $\widehat{\mathcal{B}}^k$.*

If we take a limit as $\epsilon \to 0$ (but with $r > 0$ fixed), then it is easy to see by Lemma 7.11 that (by passing along a subsequence if necessary and recoupling the laws using the Skorohod embedding theorem) we get a limiting surface in which the quantum length of $\partial\Upsilon^k_\tau$ differs from the quantum length of $\partial\Upsilon^k_\xi$ exactly by r. Moreover, we can couple this surface so that it is "close" to the original surface outside of $\mathcal{R}$ in the sense that the distortion of φ^k is small. In fact, if we take a limit as $k \to \infty$ and $c \to 0$ so that the size of $\mathcal{R}$ tends to 0 we have that φ^k converges locally uniformly to a map which is conformal everywhere except at possibly a point on $\partial\Gamma_\tau$ hence is conformal everywhere (since the map has at most an isolated singularity). The only way that the two limiting surfaces can differ is if there is an atom in the quantum area measure at this limiting point. We are now going to show that such an atom cannot form if we take a limit as $r \to 0$ simultaneously with the limit as $k \to \infty$.

Lemma 7.12. *Fix $y_0 > 0$. For each $\delta > 0$, let T_δ be uniformly random in $[\delta, 2\delta]$ and let R_δ, S_δ be random variables such that for each $\epsilon > 0$ there exists $a_0, \delta_0 > 0$ such that $a \geq a_0$ and $\delta \in (0, \delta_0)$ implies that $\mathbf{P}[X_\delta \geq a\delta^{2/3}] \leq \epsilon$ for $X_\delta \in \{R_\delta, S_\delta\}$. Suppose that Y is a $u^{3/2}$-CSBP with $Y_0 = y_0$ conditioned so that $Y_{T_\delta} = y_0 + R_\delta$ and $Y_{2\delta} = y_0 + S_\delta$. Suppose that we associate with each jump of $Y|_{[0,2\delta]}$ a conditionally independent quantum disk whose boundary length is equal to the size of the jump and let M_δ be the total amount of quantum mass of these quantum disks. Then we have that $M_\delta \to 0$ in probability as $\delta \to 0$.*

To use Lemma 7.12 to show that an atom does not form in the $k \to \infty$ limit above, note that Y plays the role of the time-reversal of the boundary length process for $\Gamma^k|_{[0,\tau]}$ in the time-interval $[\xi, \xi + 2\delta_k]$.

Proof of Lemma 7.12. The result will be a consequence of the following three observations.

First, the corresponding result holds in the case that we do not perform any conditioning on Y and the same also holds if we consider the time-reversal of Y in place of Y.

Second, if we have a $u^{3/2}$-CSBP and we condition on its value at time $t > 0$, then the law of the CSBP restricted to the time-interval $[0, t/2]$ is absolutely continuous with respect to its unconditioned law. This can be seen using a Bayes' rule calculation. The same is likewise true for the time-reversal of $Y|_{[0,t]}$ restricted to the time-interval $[t/2, t]$.



Figure 7.4: (Continuation of Figure 7.3.) Illustration of the proof of Lemma 7.10. We take $\Gamma^{1,k}|_{[0,\tau^1]}$ (resp. $\overline{\Gamma}^{1,k}|_{[0,\overline{\tau}^1]}$) to be the same as $\Gamma^{2,k}|_{[0,\sigma^2]}$ (resp. $\overline{\Gamma}^{2,k}|_{[0,\overline{\sigma}^2]}$) as path-decorated quantum surfaces. By resampling the upper left surface in $\mathcal{R}^1$ (union of yellow and green regions), we can condition so that the quantum boundary length of $\Upsilon^{1,k}$ is the same at times $\xi^1 = \widehat{\xi}^1$ and $\tau^1 = \widehat{\tau}^1$. By subsequently resampling the top right surface in $\mathcal{R}^2$, we can condition so that the quantum boundary lengths of $\widehat{\mathcal{B}}^1$ and $\widehat{\mathcal{B}}^2$ are the same. Since the tip of $\Upsilon^{1,k}$ at time $\widehat{\xi}^1$ is close in total variation to being uniformly random on $\partial\Upsilon^{1,k}$ at time $\widehat{\xi}^1$ according to the quantum boundary measure and the same is true for the tip of $\Upsilon^{2,k}$ at time $\delta_k\lfloor\sigma/\delta_k\rfloor$ (both tips shown as black squares), we can with high probability arrange so that the maps $\widehat{\phi}^k$ (resp. $\widehat{\varphi}^k$) which take $\Upsilon^{1,k}_{\widehat{\xi}^1}$ to $\Upsilon^{2,k}$ stopped at time $\delta_k\lfloor\sigma/\delta_k\rfloor$ (resp. $\widehat{\mathcal{B}}^1$ to $\widehat{\mathcal{B}}^2$) agree along $\partial\Upsilon^{1,k}\setminus\widehat{\mathcal{R}}^1$ hence, by removability, together define a conformal transformation $\widehat{\psi}^k$ in the complement of $\widehat{\mathcal{R}}^1\cup\overline{\Upsilon}^{1,k}$. By passing to a further subsequence (and recoupling) if necessary, $\widehat{\psi}^k$ converges in the limit as $k\to\infty$ to a map which is conformal in the complement of $\overline{\Gamma}^1|_{[0,\overline{\tau}^1]}$, preserves quantum area and boundary length on $\Gamma^1|_{[0,\tau^1]}$, and takes $\Gamma^1|_{[0,\tau^1]}$ to $\Gamma^2|_{[0,\sigma^2]}$.

Third, a $u^{3/2}$-CSBP run for time δ typically has fluctuations of order $\delta^{2/3}$. This follows because a 3/2-stable Lévy process run for time δ has fluctuations of order $\delta^{2/3}$ and a $u^{3/2}$-CSBP run for a short amount of time can be compared to a linear time-change of a 3/2-stable Lévy process where the speed of the time change depends on y_0; recall (2.6) and (2.7).

Combining the second and third observations with our assumption on the size of R_δ, S_δ gives that the Radon-Nikodym derivative between the conditioned and unconditioned processes does not degenerate as $\delta \to 0$. Therefore the result is a consequence of the first observation. □

We are now going to make use of Lemma 7.12 and the observations preceding it to complete the proof of Lemma 7.10; see Figure 7.4 for an illustration of the argument.

Proof of Lemma 7.10. We begin by starting off with copies $(\Gamma^{1,k}|_{[0,\tau^1]}, \overline{\Gamma}^{1,k}|_{[0,\overline{\tau}^1]}, \mathcal{B}^{1,k})$ and $(\Gamma^{2,k}|_{[0,\sigma^2]}, \overline{\Gamma}^{2,k}|_{[0,\overline{\sigma}^2]}, \overline{\mathcal{B}}^{2,k})$ respectively of $(\Gamma^k|_{[0,\tau]}, \overline{\Gamma}^k|_{[0,\overline{\tau}]}, \mathcal{B}^k)$ sampled from $\Theta_k^{x\to y}$ and $(\Gamma^k|_{[0,\sigma]}, \overline{\Gamma}^k|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}}^k)$ sampled from $\Theta_k^{y\to x}$. We take these to be coupled together so that (as path-decorated quantum surfaces) we have $\Gamma^{1,k}|_{[0,\tau^1]} = \Gamma^{2,k}|_{[0,\sigma^2]}$, $\overline{\Gamma}^{1,k}|_{[0,\overline{\tau}^1]} = \overline{\Gamma}^{2,k}|_{[0,\overline{\sigma}^2]}$, and $\mathcal{B}^{1,k} = \overline{\mathcal{B}}^{2,k}$. Let $\xi^j < \tau^j < \zeta^j$ for $j = 1, 2$ be as in Lemma 7.11 for the two triples and let $\mathcal{R}^j$ and E_j respectively be the corresponding surfaces and events. We next claim that the probability that both E_1 and E_2 occur tends to 1 as $k \to \infty$ and then $c \to 0$. Indeed, this follows because we can think of first generating the QLE(8/3, 0) process which corresponds to $\Gamma^{1,k}$ up until time ζ^1 before drawing $\overline{\Gamma}^{1,k}$. Then E_1 is the event that $\overline{\Gamma}^{1,k}$ does not hit the boundary segment which corresponds to the extra $\zeta^1 - \tau^1 \in [0, c)$ units of time for the SLE_6 starting from the tip of $\Gamma^{1,k}|_{[0,\tau^1]}$ before hitting $\partial\Gamma^{1,k}_{\tau^1}$. Arguing as in Lemma 5.4, the probability of hitting this segment is proportional to its quantum length. Hence the claim follows because it is easy to see that this quantum length tends to 0 in probability as $k \to \infty$ and then $c \to 0$. The same argument implies that the probability that E_2 occurs tends to 1 as $k \to \infty$ and then $c \to 0$.

We apply Lemma 7.11 as explained just above to obtain surfaces $(\Upsilon^{1,k}, \overline{\Upsilon}^{1,k}, \widehat{\mathcal{B}}^{1,k})$ and $(\Upsilon^{2,k}, \overline{\Upsilon}^{2,k}, \widehat{\mathcal{B}}^{2,k})$ as follows:

1. We resample the surface inside of $\mathcal{R}^1$ (with the quantum length of $\partial\mathcal{R}^1$ fixed) so that the quantum boundary length of $\partial\Upsilon^{1,k}$ is the same at the times $\widehat{\xi}^1 = \xi^1$ and $\widehat{\tau}^1 = \tau^1$

2. We then resample the surface in $\mathcal{R}^2$ (with the quantum length of $\partial\mathcal{R}^2$ fixed) so that the change in the quantum boundary length of $\partial\Upsilon^{2,k}$ from time $\widehat{\xi}^2 = \xi^2$ to time $\widehat{\tau}^2 = \tau^2$ is minus the change in the quantum boundary length of $\partial\Gamma^{1,k}_{\xi^1}$ to $\partial\Gamma^{1,k}_{\tau^1}$.

The two changes to the quantum boundary length cancel so that the quantum boundary length of $\widehat{\mathcal{B}}^{1,k}$ is the same as the quantum boundary length of $\widehat{\mathcal{B}}^{2,k}$.

We note that the conditional law of each of the $\widehat{\mathcal{B}}^{j,k}$ given its quantum boundary length is not exactly that of a quantum disk. This is because the conditional law of the quantum boundary length of $\partial(\widehat{\mathcal{R}}^1 \cup \Upsilon^{1,k} \cup \overline{\Upsilon}^{1,k})$ given the quantum boundary length of $\partial(\Upsilon^{1,k} \cup \overline{\Upsilon}^{1,k})$ is not the same as the conditional law of the quantum boundary length of $\partial(\mathcal{R}^1 \cup \Gamma^1_\tau \cup \overline{\Gamma}^1_{\overline{\tau}})$ given the quantum boundary length of $\partial(\Gamma^1_\tau \cup \overline{\Gamma}^1_{\overline{\tau}})$. However, it is not difficult to see that the total variation distance between the law of the surface and that of a quantum disk with the given boundary length can be made arbitrarily small by taking k to be sufficiently large with $c > 0$ small fixed. The same is likewise true with $\widehat{\mathcal{B}}^{2,k}$ in place of $\widehat{\mathcal{B}}^{1,k}$. Therefore, by choosing first $c > 0$ small and then k sufficiently large, we can couple each of the $\widehat{\mathcal{B}}^{j,k}$ to be the same with high probability (note that both surfaces have the same boundary length). That is, with high probability there exists a conformal map $\widehat{\varphi}^k$ which takes $\widehat{\mathcal{B}}^{1,k}$ to $\widehat{\mathcal{B}}^{2,k}$ and preserves both quantum boundary length and area. Similarly, there is also a conformal map $\widehat{\phi}^k$ which takes $\Upsilon^{1,k}_{\widehat{\xi}^1}$ to $\Upsilon^{2,k}$ stopped at time $\delta_k \lfloor \sigma/\delta_k \rfloor$ and preserves both quantum boundary length and area. By the construction of QLE$(8/3, 0)$, we know that the law of the tip of $\Upsilon^{1,k}$ at time $\widehat{\xi}^1$ is in total variation close to being uniformly distributed according to quantum boundary measure on $\partial\Upsilon^{1,k}$ and the same is likewise true for the tip of $\Upsilon^{2,k}$ at time $\delta_k \lfloor \sigma/\delta_k \rfloor$. Therefore we can arrange so that $\widehat{\varphi}^k$ also takes the tip of $\Upsilon^{1,k}$ at time $\widehat{\xi}^1$ to the tip of $\Upsilon^{2,k}$ at time $\delta_k \lfloor \sigma/\delta_k \rfloor$ with probability tending to 1 as $k \to \infty$.

On this high probability event, we therefore have that $\widehat{\varphi}^k$ and $\widehat{\phi}^k$ agree on $\partial\Upsilon^{1,k} \setminus \mathcal{R}^1$. (This follows because the quantum boundary length of $\partial\Upsilon^{1,k}$ is the same as time $\widehat{\xi}^1$ and at time $\widehat{\tau}^1$.) That is, the map $\widehat{\psi}^k$ which takes $(\Upsilon^{1,k} \cup \widehat{\mathcal{B}}^{1,k}) \setminus \widehat{\mathcal{R}}^1$ to $\Upsilon^{2,k} \cup \widehat{\mathcal{B}}^{2,k}$ given by $\widehat{\varphi}^k$ on $\widehat{\mathcal{B}}^{1,k} \setminus \widehat{\mathcal{R}}^1$ and $\widehat{\phi}^k$ on $\Upsilon^{1,k} \setminus \widehat{\mathcal{R}}^1$ is conformal on the interiors of $\widehat{\mathcal{B}}^{1,k} \setminus \widehat{\mathcal{R}}^1$ and $\Upsilon^{1,k} \setminus \widehat{\mathcal{R}}^1$ and a homeomorphism on its entire domain. Therefore by the conformal removability of $\partial\Upsilon^{1,k}_{\widehat{\xi}^1}$, we have that $\widehat{\psi}^k$ is conformal on the interior of its domain.

Since $\operatorname{diam}(\widehat{\mathcal{R}}^1) \to 0$ in probability as $k \to \infty$, it therefore follows that, by passing to a further subsequence if necessary (and recoupling the laws using the Skorohod embedding theorem), $\widehat{\psi}^k$ converges almost surely as $k \to \infty$ with respect to the local uniform topology to a map $\widehat{\psi} \colon \Gamma^1_{\tau^1} \cup \mathcal{B}^1 \to \Gamma^2_{\sigma^2} \cup \overline{\mathcal{B}}^2$ which is a homeomorphism and conformal in its interior. By Lemma 7.6, the interior of the domain of $\widehat{\psi}$ includes all of $\Gamma^1_{\tau^1}$ except possibly a single point — the unique point where it hits $\overline{\Gamma}|^1_{[0,\overline{\tau}^1]}$. However, it is not difficult to see that $\widehat{\psi}$ is continuous at this point because $\widehat{\varphi}^k$ is continuous at the unique point where $\Gamma|^{1,k}_{[0,\tau^1]}$ hits $\overline{\Gamma}|^{1,k}_{[0,\overline{\tau}^1]}$ uniformly in k. It therefore follows that the distribution of $\Gamma|_{[0,\tau]}$ under $\Theta^{x\to y}$ is the same as that of $\Gamma|_{[0,\sigma]}$ under $\Theta^{y\to x}$ (in the sense described just before the statement of the lemma). The same argument also implies that the distribution of $\overline{\Gamma}|_{[0,\overline{\tau}]}$ under $\Theta^{x\to y}$ is the same as the distribution of $\overline{\Gamma}|_{[0,\overline{\sigma}]}$ under $\Theta^{y\to x}$. The result follows because $\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B}$ are conditionally independent under

$\Theta^{x\to y}$ given their boundary lengths, the same holds under $\Theta^{y\to x}$, and the joint law of the boundary lengths is the same as that of $\Gamma|_{[0,\sigma]}$, $\overline{\Gamma}|_{[0,\overline{\sigma}]}$, $\overline{\mathcal{B}}$ under $\Theta^{y\to x}$. □

7.2.4 Proof of Theorem 7.1

Lemma 7.10 implies that the $\Theta^{x\to y}$ distribution of $(\Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]}, \mathcal{B})$ is equal to the $\Theta^{y\to x}$ distribution of $(\Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]}, \overline{\mathcal{B}})$ (each viewed as a quantum surface in the sense described just before the statement of Lemma 7.8). This implies that we can construct a coupling of copies $(\Gamma^1|_{[0,\tau^1]}, \overline{\Gamma}^1|_{[0,\overline{\tau}^1]}, \mathcal{B}^1)$ and $(\Gamma^2|_{[0,\sigma^2]}, \overline{\Gamma}^2|_{[0,\overline{\sigma}^2]}, \overline{\mathcal{B}}^2)$ so that (as quantum surfaces in the sense described just before the statement of Lemma 7.8) we have $\Gamma^1_{\tau^1} = \Gamma^2_{\sigma^2}$, $\overline{\Gamma}^1_{\overline{\tau}^1} = \overline{\Gamma}^2_{\overline{\sigma}^2}$, and $\mathcal{B}^1 = \overline{\mathcal{B}}^2$. This implies the existence of homeomorphisms $\varphi\colon \Gamma^1_{\tau^1} \to \Gamma^2_{\sigma^2}$, $\overline{\varphi}\colon \overline{\Gamma}^1_{\overline{\tau}^1} \to \overline{\Gamma}^2_{\overline{\sigma}^2}$, and $\psi\colon \mathcal{B}^1 \to \overline{\mathcal{B}}^2$ which are each conformal in the interior of their domain and preserve both quantum boundary length. By Lemma 7.7, we know that the point on $\partial\Gamma^1_{\tau^1}$ which is glued to the almost surely unique pinch point on $\partial\mathcal{B}^1$ is uniformly random from the quantum boundary measure on $\partial\Gamma^1_{\tau^1}$ and the same is likewise true for $\Gamma^2_{\sigma^2}$. Thus we can couple so that φ (resp. $\overline{\varphi}$) sends this point to the corresponding point on $\partial\Gamma^2_{\sigma^2}$ (resp. $\partial\overline{\Gamma}^2_{\overline{\sigma}^2}$). Let $\mathcal{S}^j$ for $j = 1, 2$ be the doubly-marked quantum sphere associated with $(\Gamma^1_{\tau^1}, \overline{\Gamma}^1_{\overline{\tau}^1}, \mathcal{B}^1)$ and $(\Gamma^2_{\sigma^2}, \overline{\Gamma}^2_{\overline{\sigma}^2}, \overline{\mathcal{B}}^2)$, respectively. Letting ϕ be the map $\mathcal{S}^1 \to \mathcal{S}^2$ which on $\Gamma^1_{\tau^1}$ (resp. $\overline{\Gamma}^1_{\overline{\tau}^1}$) is given by φ (resp. $\overline{\varphi}$) and on $\mathcal{B}^1$ is given by ψ, we thus see that ϕ is a homeomorphism which is conformal in the complement of $\partial\mathcal{B}^1 = \partial\Gamma^1_{\tau^1} \cup \partial\overline{\Gamma}^1_{\overline{\tau}^1}$. By the conformal removability of $\partial\mathcal{B}^1$ (Lemma 7.6), we thus have that ϕ is conformal everywhere hence must be a Möbius transformation, which completes the proof. □

7.2.5 Conditional law of the unexplored surface

We finish this section by stating the analog of the final part of Proposition 4.1 for QLE$(8/3, 0)$.

Proposition 7.13. *The $\Theta^{x\to y}$ conditional distribution given $(\mathcal{S}, x, y)$ and $\Gamma|_{[0,\tau]}$ is equal to the Θ conditional distribution given $(\mathcal{S}, x, y)$ and $\Gamma|_{[0,\tau]}$.*

Proof. The last part of Proposition 4.1 implies that the analogous statement holds for each of the δ-approximations to QLE$(8/3, 0)$. Therefore it follows that this property holds in the limit. □

8 Metric construction

The purpose of this section is to show that QLE$(8/3, 0)$ equipped with the quantum distance parameterization defines a metric on a countable, dense subset consisting of

a sequence of i.i.d. points chosen from the quantum measure of a $\sqrt{8/3}$-LQG sphere. The approach we take, as explained and outlined in Section 1.1, is based on ideas from a joint work by the second co-author, Sam Watson, and Hao Wu.

We begin by considering a pair of QLE$(8/3, 0)$ explorations Γ and $\overline{\Gamma}$ on a doubly marked quantum sphere $(\mathcal{S}, x, y)$ where the first (resp. second) process starts from x (resp. y) and is targeted at y (resp. x). We then use several results accumulated earlier in the paper to prove that the "distances" computed by these two explorations are almost surely the same.

The remainder of this section is structured as follows. In Section 8.1, we describe the setup and notation that we will use throughout the rest of the section. We will complete the proof of Theorem 1.1 in Section 8.2.

8.1 Setup and notation

We will use the same setup and notation described in the beginning of Section 7. Since $\Theta^{x\to y}$ and $\Theta^{y\to x}$ are infinite measures, we cannot normalize them to be probability measures. However, these measures conditioned on certain quantities do yield probability measures. In these cases, we will let $\mathbf{P}_{\Theta^{x\to y}}[\cdot \,|\, \cdot]$ (resp. $\mathbf{P}_{\Theta^{y\to x}}[\cdot \,|\, \cdot]$) and $\mathbf{E}_{\Theta^{x\to y}}[\cdot \,|\, \cdot]$ (resp. $\mathbf{E}_{\Theta^{y\to x}}[\cdot \,|\, \cdot]$) denote the corresponding probability and expectation. We will similarly write $\mathbf{P}_{\Theta}[\cdot \,|\, \cdot]$ and $\mathbf{E}_{\Theta}[\cdot \,|\, \cdot]$ for the probability and expectation which correspond to Θ conditioned on a quantity which leads to a probability measure. For each $t \geq 0$, we also let $\mathcal{F}_t = \sigma(\mathcal{S}, x, y, \Gamma_s : s \leq t)$ and $\overline{\mathcal{F}}_t = \sigma(\mathcal{S}, x, y, \overline{\Gamma}_s : s \leq t)$. That is, $\mathcal{F}_t$ (resp. $\overline{\mathcal{F}}_t$) is the filtration generated by $(\mathcal{S}, x, y)$ and Γ (resp. $\overline{\Gamma}$) with the quantum distance parameterization.

8.2 Coupling explorations and the metric property

We will complete the proof of Theorem 1.1 in this section. The first step is to prove the following slightly stronger form of Lemma 1.2, which is in some sense the heart of the matter.

Proposition 8.1. *The $\Theta^{x\to y}$ distribution of $(\mathcal{S}, x, y, \tau, \Gamma, \overline{\tau}, \overline{\Gamma})$ is the same as the $\Theta^{y\to x}$ distribution of $(\mathcal{S}, x, y, \sigma, \Gamma, \overline{\sigma}, \overline{\Gamma})$.*

Theorem 7.1 implies that the $\Theta^{x\to y}$ distribution of $(\mathcal{S}, x, y, \Gamma|_{[0,\tau]}, \overline{\Gamma}|_{[0,\overline{\tau}]})$ is equal to the $\Theta^{y\to x}$ distribution of $(\mathcal{S}, x, y, \Gamma|_{[0,\sigma]}, \overline{\Gamma}|_{[0,\overline{\sigma}]})$. To prove Proposition 8.1, we will try to understand the conditional law of Γ and $\overline{\Gamma}$ given this information. Our argument will follow the sketch given at the end of Section 1.1.

We begin by fixing a bounded function F and let

$$M_t = \mathbf{E}_{\Theta}[F(\Gamma) \,|\, \mathcal{F}_t] \quad \text{and} \quad \overline{M}_t = \mathbf{E}_{\Theta}[F(\overline{\Gamma}) \,|\, \overline{\mathcal{F}}_t] \quad \text{for each} \quad t \geq 0.$$

Since M_t and $\overline{M}_t$ are both continuous time, bounded martingales, we note that they both a.s. jump at most a countable number of times. This property will be important for us below. The following lemma is the main input into the proof of Proposition 8.1.

Lemma 8.2. *We have on a set of full Θ measure both*

$$M_\sigma = \mathbf{E}_{\Theta^{y\to x}}[F(\Gamma)\,|\,\mathcal{F}_\sigma, \overline{\mathcal{F}}_{\overline{\sigma}}] \quad \textit{and} \quad \overline{M}_{\overline{\tau}} = \mathbf{E}_{\Theta^{x\to y}}[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_{\overline{\tau}}]. \tag{8.1}$$

Proof. We will only establish the second equality in (8.1); the proof of the first equality is analogous. We first recall that on a set of full Θ measure we have $\mathbf{E}_{\Theta^{x\to y}}[\cdot\,|\,\mathcal{F}_\tau] = \mathbf{E}_\Theta[\cdot\,|\,\mathcal{F}_\tau]$; see Proposition 7.13. Throughout, we will write dt for Lebesgue measure on $\mathbf{R}_+$. We now observe that $d\Theta \otimes dt$ almost everywhere we have that

$$\mathbf{E}_\Theta[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_t] = \mathbf{E}_{\Theta^{x\to y}}[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_t]. \tag{8.2}$$

Since $\mathcal{F}_\tau$ and $\overline{\mathcal{F}}_t$ are conditionally independent given $\mathcal{S}$ under Θ, we have $d\Theta \otimes dt$ almost everywhere that

$$\overline{M}_t = \mathbf{E}_\Theta[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_t]. \tag{8.3}$$

Combining (8.2) and (8.3), we have $d\Theta \otimes dt$ almost everywhere that

$$\overline{M}_t = \mathbf{E}_{\Theta^{x\to y}}[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_t]. \tag{8.4}$$

In particular, the event E that (8.4) holds for all rational times simultaneously has full Θ measure.

We will deduce (8.1) from (8.4) by showing that on a set of full Θ measure we have both

$$\overline{M}_t \to \overline{M}_{\overline{\tau}} \quad \text{as} \quad t \to \overline{\tau},\ t < \overline{\tau},\ t \in \mathbf{Q} \quad \text{and} \tag{8.5}$$

$$\mathbf{E}_{\Theta^{x\to y}}[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_t] \to \mathbf{E}_{\Theta^{x\to y}}[F(\overline{\Gamma})\,|\,\mathcal{F}_\tau, \overline{\mathcal{F}}_{\overline{\tau}}] \quad \text{as} \quad t \to \overline{\tau},\ t < \overline{\tau},\ t \in \mathbf{Q}. \tag{8.6}$$

We emphasize that in (8.5) and (8.6) we take the limit along rational $t < \overline{\tau}$.

To prove (8.6) it suffices to show that the σ-algebra generated by $\overline{\mathcal{F}}_t$ for $t < \overline{\tau}$ is equal to $\overline{\mathcal{F}}_{\overline{\tau}}$. It in turn suffices to show that the closure of $\cup_{t<\overline{\tau}} \overline{\Gamma}_t$ is almost surely equal to $\overline{\Gamma}_{\overline{\tau}}$ which will follow by showing that $\overline{\tau}$ does not correspond to a jump in the time change from capacity to quantum distance time. Since there can only be a countable number of such jump times and $\overline{\sigma}$ is uniform in $[0, \overline{d}_{\mathcal{Q}}(y,x)]$, it follows that $\overline{\sigma}$ almost surely does not correspond to such a jump time for $\overline{\Gamma}$. By Theorem 7.1, the $\Theta^{x\to y}$ distribution of $\overline{\Gamma}|_{[0,\overline{\tau}]}$ is the same as the $\Theta^{y\to x}$ distribution of $\overline{\Gamma}|_{[0,\overline{\sigma}]}$. We therefore conclude that $\overline{\tau}$ similarly does not correspond to a time at which the capacity of $\overline{\Gamma}$ jumps.

The existence of the limit in (8.6) combined with (8.4) implies that $\overline{M}_t$ almost surely has a limit as $t \to \overline{\tau}$ along rationals $t < \overline{\tau}$. We need to show that this limit is almost surely

equal to $\overline{M}_{\overline{\tau}}$. As we remarked above, we know that $\overline{M}_t$ can jump at most countably many times as it is a continuous-time martingale. Since $\mathbf{P}_\Theta[\overline{\sigma} = t \,|\, \overline{\mathcal{F}}_{\overline{d}_{\mathcal{Q}}(x,y)}] = 0$ almost surely for any fixed t, it follows that $\overline{\sigma}$ is almost surely not a jump time for $\overline{M}_t$. Consequently, we have that

$$\lim_{t \to \overline{\sigma}^-} \overline{M}_t = \overline{M}_{\overline{\sigma}} \tag{8.7}$$

on a set of full Θ measure. By Theorem 7.1, we know that the $\Theta^{y \to x}$ distribution of $\overline{M}|_{[0,\overline{\sigma}]}$ is equal to the $\Theta^{x \to y}$ distribution of $\overline{M}|_{[0,\overline{\tau}]}$. Therefore (8.7) implies that $\lim_{t \to \overline{\tau}^-} \overline{M}_t = \overline{M}_{\overline{\tau}}$ on a set of full $\Theta^{x \to y}$ measure. Since Θ is absolutely continuous with respect to $\Theta^{x \to y}$, it therefore follows that $\lim_{t \to \overline{\tau}^-} \overline{M}_t = \overline{M}_{\overline{\tau}}$ on a set of full Θ measure. Combining everything completes the proof. □

Proof of Proposition 8.1. Theorem 7.1 implies that the $\Theta^{x \to y}$ distribution of $(\mathcal{S}, x, y)$, $\Gamma|_{[0,\tau]}$, and $\overline{\Gamma}|_{[0,\overline{\tau}]}$ is equal to the $\Theta^{y \to x}$ distribution of $(\mathcal{S}, x, y)$, $\Gamma|_{[0,\sigma]}$, and $\overline{\Gamma}|_{[0,\overline{\sigma}]}$.

We claim that the $\Theta^{x \to y}$ conditional law of Γ given $\mathcal{F}_\tau, \overline{\mathcal{F}}_{\overline{\tau}}$ is the same as the $\Theta^{y \to x}$ conditional law of Γ given $\mathcal{F}_\sigma, \overline{\mathcal{F}}_{\overline{\sigma}}$. The same argument will give that the $\Theta^{y \to x}$ conditional law of $\overline{\Gamma}$ given $\mathcal{F}_\sigma, \overline{\mathcal{F}}_{\overline{\sigma}}$ is the same as the $\Theta^{x \to y}$ conditional law of $\overline{\Gamma}$ given $\mathcal{F}_\tau, \overline{\mathcal{F}}_{\overline{\tau}}$. Upon showing this, the proof will be complete. To see the claim, we first note that Proposition 7.13 implies that the $\Theta^{x \to y}$ conditional law of Γ given $\mathcal{F}_\tau, \overline{\mathcal{F}}_{\overline{\tau}}$ is the same as its Θ conditional law given $\mathcal{F}_\tau, \overline{\mathcal{F}}_{\overline{\tau}}$. By the Θ conditional independence of $\mathcal{F}_\tau$ and $\overline{\mathcal{F}}_{\overline{\tau}}$ given $(\mathcal{S}, x, y)$, this conditional law is in turn equal to the Θ conditional law of Γ given $\mathcal{F}_\tau$. Lemma 8.2 then implies that the Θ conditional law of Γ given $\mathcal{F}_\tau$ is equal to the $\Theta^{y \to x}$ conditional law of Γ given $\mathcal{F}_\sigma, \overline{\mathcal{F}}_{\overline{\sigma}}$ (this follows from the first equation in (8.1)). This proves the claim, hence the proposition. □

Proposition 8.3. *We have on a set of full Θ measure that $d_{\mathcal{Q}}(x, y) = \overline{d}_{\mathcal{Q}}(y, x)$ and that the common value of $d_{\mathcal{Q}}(x, y)$ and $\overline{d}_{\mathcal{Q}}(y, x)$ is almost surely determined by $(\mathcal{S}, x, y)$.*

Proof. Proposition 8.1 implies that

$$d_{\mathcal{Q}}(x, y) = \frac{d\Theta^{x \to y}}{d\Theta} = \frac{d\Theta^{y \to x}}{d\Theta} = \overline{d}_{\mathcal{Q}}(y, x).$$

This implies that $d_{\mathcal{Q}}(x, y) = \overline{d}_{\mathcal{Q}}(y, x)$ on a set of full Θ measure. Moreover, the common value of $d_{\mathcal{Q}}(x, y)$ and $\overline{d}_{\mathcal{Q}}(y, x)$ is almost surely determined by $(\mathcal{S}, x, y)$ because we took Γ and $\overline{\Gamma}$ to be conditionally independent given $(\mathcal{S}, x, y)$ under Θ. In particular, this implies that $d_{\mathcal{Q}}(x, y)$ and $\overline{d}_{\mathcal{Q}}(y, x)$ are conditionally independent given $(\mathcal{S}, x, y)$ and the only way that two conditionally independent random variables given $(\mathcal{S}, x, y)$ can be equal on a set of full measure is if they are almost surely determined by $(\mathcal{S}, x, y)$. □

Proposition 8.1 implies that if $\mathcal{S}$ has the law of a unit area quantum sphere and $x, y \in \mathcal{S}$ are chosen independently from the quantum measure on $\mathcal{S}$, then the amount of time that it takes a QLE$(8/3, 0)$ with the quantum distance parameterization starting from

x to reach y is almost surely determined by $\mathcal{S}$ and is equal to the amount of time that it takes for a QLE$(8/3, 0)$ with the quantum distance parameterization starting from y to reach x. Moreover, this quantity does not depend on the choice of sequence that we chose in the construction of the QLE$(8/3, 0)$. Therefore from now on we will only write $d_{\mathcal{Q}}$ (and not $\overline{d}_{\mathcal{Q}}$). Conditionally on $\mathcal{S}$, we let (x_n) be a sequence of i.i.d. points picked from the quantum area measure. For each i, j, we let $d_{\mathcal{Q}}(x_i, x_j)$ be the amount of quantum distance time that it takes for a QLE$(8/3, 0)$ starting from x_i to reach x_j. Then it follows that $d_{\mathcal{Q}}(x_i, x_j) = d_{\mathcal{Q}}(x_j, x_i)$ for all i, j and $d_{\mathcal{Q}}$ is almost surely determined by $\mathcal{S}$. Moreover, it is immediate from the construction that $d_{\mathcal{Q}}(x_i, x_j) > 0$ almost surely for any $i \neq j$.

Our next goal is to establish Proposition 8.5, which will be used in the proof of Theorem 1.1 to show that $d_{\mathcal{Q}}$ satisfies the triangle inequality hence is a distance function on (x_n). Before we state and prove this result, we need to collect the following elementary lemma.

Lemma 8.4. *Fix $D > 0$. Suppose that $F\colon [0, D] \to [0, D]$ is a non-increasing function such that if U is uniform in $[0, D]$ then $F(U)$ is uniform in $[0, D]$. Then $F(d) = D - d$ for all $d \in [0, D]$.*

Proof. This is essentially obvious, but let us explain the point to be clear. Since $F(U)$ is uniform in $[0, D]$, we have $\mathbf{P}[F(U) \geq d] = 1 - d/D$ for all $d \in [0, D]$. Since F is non-increasing and $F(U)$ is uniform it follows that there cannot be a non-empty open interval in $[0, D]$ on which F is constant. Consequently, $\mathbf{P}[F(U) \leq F(d)] = \mathbf{P}[U \geq d] = 1 - d/D$. Since $V = F(U)$ is uniform and we have shown that $\mathbf{P}[V \leq F(d)] = 1 - d/D$ for all $d \in [0, D]$, we conclude that $F(d) = D - d$ for all $d \in [0, D]$. □

Proposition 8.5. *On a set of full Θ measure, we have that $\tau + \overline{\tau} = D$ where D is the common value of $d_{\mathcal{Q}}(x, y)$ and $d_{\mathcal{Q}}(y, x)$.*

Proof. It suffices to show that $\tau + \overline{\tau} = D$ under $\Theta^{x \to y}$ since this measure is mutually absolutely continuous with respect to Θ.

For each $d \in [0, D]$, let $F(d)$ be the first time that $\overline{\Gamma}$ hits Γ_d. We emphasize that F is determined by Γ, $\overline{\Gamma}$, and $\mathcal{S}$. We already know that $d_{\mathcal{Q}}(x, y) = d_{\mathcal{Q}}(y, x)$ a.s. and that this quantity is a.s. determined by $(\mathcal{S}, x, y)$. We know that $U = \tau / d_{\mathcal{Q}}(x, y)$ is uniform in $[0, 1]$ conditionally on $(\mathcal{S}, x, y)$ and Γ. The symmetry we have established in Proposition 8.1 then implies that $\overline{U} = \overline{\tau} / d_{\mathcal{Q}}(y, x)$ is uniform in $[0, 1]$ conditionally on $(\mathcal{S}, x, y)$ and $\overline{\Gamma}$. In the case of τ, it is clearly also uniform in $[0, D]$ once we condition on $(\mathcal{S}, x, y, \Gamma, \overline{\Gamma})$, which determines F. That this is true for $\overline{\tau}$ as well follows from Proposition 8.1, so that both τ and $\overline{\tau}$ are uniform in $[0, D]$ conditionally on F. To finish the proof of the proposition, it suffices to show that $F(d) = D - d$. We know that $F\colon [0, D] \to [0, D]$ is non-decreasing and that $F(\tau) = \overline{\tau}$. Therefore the result follows from Lemma 8.4. □

Proof of Theorem 1.1. Let $d_{\mathcal{Q}}$ and (x_n) be as defined just before the statement of Lemma 8.4. To finish the proof, it suffices to show that $d_{\mathcal{Q}}$ a.s. satisfies the strict triangle inequality. Suppose that $x, y, z \in (x_n)$ are distinct. We will argue that a.s. we have

$$d_{\mathcal{Q}}(x, z) < d_{\mathcal{Q}}(x, y) + d_{\mathcal{Q}}(y, z). \tag{8.8}$$

We shall assume that we are working on the event that $d_{\mathcal{Q}}(x, y) < d_{\mathcal{Q}}(x, z)$, for otherwise (8.8) is trivial. Consider the QLE$(8/3, 0)$ growth Υ starting from x and stopped upon hitting y. Given this, we then consider the QLE$(8/3, 0)$ growth $\overline{\Upsilon}$ starting from z and stopped upon hitting Υ. Then we must have that the radius of this growth is at most $d_{\mathcal{Q}}(y, z)$ because $y \in \Upsilon$. As it is easy to see that the a.s. unique point in $\overline{\Upsilon} \cap \Upsilon$ is uniformly distributed in $\partial\Upsilon$ according to the quantum measure, we a.s. have that $y \notin \overline{\Upsilon}$. That is, the radius of $\overline{\Upsilon}$ a.s. is strictly less than $d_{\mathcal{Q}}(y, z)$. On the other hand, Proposition 8.5 implies that the sum of $d_{\mathcal{Q}}(x, y)$ and the radius of this growth is a.s. equal to $d_{\mathcal{Q}}(x, z)$. Combining proves (8.8), which completes the proof. □

References

[AHS15] J. Aru, Y. Huang, and X. Sun. Two perspectives of the unit area quantum sphere and their equivalence. *ArXiv e-prints*, December 2015, 1512.06190.

[Ber96] J. Bertoin. *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996. MR1406564 (98e:60117)

[Bis07] C. J. Bishop. Conformal welding and Koebe's theorem. *Ann. of Math. (2)*, 166(3):613–656, 2007. MR2373370 (2009d:30010)

[CL12] N. Curien and J.-F. Le Gall. The Brownian plane. *ArXiv e-prints*, April 2012, 1204.5921.

[CL14a] N. Curien and J.-F. Le Gall. Scaling limits for the peeling process on random maps. *ArXiv e-prints*, December 2014, 1412.5509.

[CL14b] N. Curien and J.-F. Le Gall. The hull process of the Brownian plane. *ArXiv e-prints*, September 2014, 1409.4026.

[DKRV14] F. David, A. Kupiainen, R. Rhodes, and V. Vargas. Liouville Quantum Gravity on the Riemann sphere. *ArXiv e-prints*, October 2014, 1410.7318.

[DMS14] B. Duplantier, J. Miller, and S. Sheffield. Liouville quantum gravity as a mating of trees. *ArXiv e-prints*, September 2014, 1409.7055.

[DRV15] F. David, R. Rhodes, and V. Vargas. Liouville Quantum Gravity on the complex tori. *ArXiv e-prints*, April 2015, 1504.00625.

[DS11] B. Duplantier and S. Sheffield. Liouville quantum gravity and KPZ. *Invent. Math.*, 185(2):333–393, 2011. 0808.1560. MR2819163 (2012f:81251)

[GM16] E. Gwynne and J. Miller. Convergence of the topology of critical Fortuin-Kasteleyn planar maps to that of CLE_κ on a Liouville quantum surface. 2016. In preparation.

[GS15a] E. Gwynne and X. Sun. Scaling limits for the critical Fortuin-Kastelyn model on a random planar map II: local estimates and empty reduced word exponent. *ArXiv e-prints*, May 2015, 1505.03375.

[GS15b] E. Gwynne and X. Sun. Scaling limits for the critical Fortuin-Kastelyn model on a random planar map III: finite volume case. *ArXiv e-prints*, October 2015, 1510.06346.

[JS00] P. W. Jones and S. K. Smirnov. Removability theorems for Sobolev functions and quasiconformal maps. *Ark. Mat.*, 38(2):263–279, 2000. MR1785402 (2001i:46049)

[Kri05] M. Krikun. Uniform infinite planar triangulation and related time-reversed critical branching process. *Journal of Mathematical Sciences*, 131(2):5520–5537, 2005.

[Kyp06] A. E. Kyprianou. *Introductory lectures on fluctuations of Lévy processes with applications*. Universitext. Springer-Verlag, Berlin, 2006. MR2250061 (2008a:60003)

[Lam67] J. Lamperti. Continuous state branching processes. *Bull. Amer. Math. Soc.*, 73:382–386, 1967. MR0208685 (34 #8494)

[LG99] J.-F. Le Gall. *Spatial branching processes, random snakes and partial differential equations*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1999. MR1714707 (2001g:60211)

[LG13] J.-F. Le Gall. Uniqueness and universality of the Brownian map. *Ann. Probab.*, 41(4):2880–2960, 2013. 1105.4842. MR3112934

[LG14] J.-F. Le Gall. The Brownian map: a universal limit for random planar maps. In *XVIIth International Congress on Mathematical Physics*, pages 420–428. World Sci. Publ., Hackensack, NJ, 2014. MR3204495

[LGP08] J.-F. Le Gall and F. Paulin. Scaling limits of bipartite planar maps are homeomorphic to the 2-sphere. *Geom. Funct. Anal.*, 18(3):893–918, 2008. math/0612315. MR2438999 (2010a:60030)

[Mie13] G. Miermont. The Brownian map is the scaling limit of uniform random plane quadrangulations. *Acta Math.*, 210(2):319–401, 2013. MR3070569

[MM06] J.-F. Marckert and A. Mokkadem. Limit of normalized quadrangulations: the Brownian map. *Ann. Probab.*, 34(6):2144–2202, 2006. MR2294979 (2007m:60092)

[MS12a] J. Miller and S. Sheffield. Imaginary Geometry I: Interacting SLEs. *ArXiv e-prints*, January 2012, 1201.1496. To appear in Probability Theory and Related Fields.

[MS12b] J. Miller and S. Sheffield. Imaginary geometry II: reversibility of $\mathrm{SLE}_\kappa(\rho_1;\rho_2)$ for $\kappa \in (0,4)$. *ArXiv e-prints*, January 2012, 1201.1497. To appear in Annals of Probability.

[MS12c] J. Miller and S. Sheffield. Imaginary geometry III: reversibility of SLE_κ for $\kappa \in (4,8)$. *ArXiv e-prints*, January 2012, 1201.1498. To appear in Annals of Math.

[MS13a] J. Miller and S. Sheffield. Imaginary geometry IV: interior rays, whole-plane reversibility, and space-filling trees. *ArXiv e-prints*, February 2013, 1302.4738.

[MS13b] J. Miller and S. Sheffield. Quantum Loewner Evolution. *ArXiv e-prints*, December 2013, 1312.5745. To appear in Duke Mathematical Journal.

[MS15a] J. Miller and S. Sheffield. An axiomatic characterization of the Brownian map. *ArXiv e-prints*, June 2015, 1506.03806.

[MS15b] J. Miller and S. Sheffield. Liouville quantum gravity spheres as matings of finite-diameter trees. *ArXiv e-prints*, June 2015, 1506.03804.

[MS16a] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map II: geodesics and continuity of the embedding. 2016. In preparation.

[MS16b] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map III: the conformal structure is determined. 2016. In preparation.

[MW13] J. Miller and H. Wu. Intersections of SLE Paths: the double and cut point dimension of SLE. *ArXiv e-prints*, March 2013, 1303.4725.

[RS05] S. Rohde and O. Schramm. Basic properties of SLE. *Ann. of Math. (2)*, 161(2):883–924, 2005.

[She11] S. Sheffield. Quantum gravity and inventory accumulation. *ArXiv e-prints*, August 2011, 1108.2241. To appear in Annals of Probability.

[She15] S. Sheffield. Conformal weldings of random surfaces: SLE and the quantum gravity zipper. *ArXiv e-prints*, December 2015, 1012.4797. To appear in Annals of Probability.

[SS13] O. Schramm and S. Sheffield. A contour line of the continuum Gaussian free field. *Probab. Theory Related Fields*, 157(1-2):47–80, 2013. 1008.2447. MR3101840

Department of Mathematics
Massachusetts Institute of Technology
Cambridge, MA, USA